


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C07D 207/09, A61K 31/40, C07D 401/12, 401/14, 403/12, 403/14, 405/06, 405/12, 405/14, 409/06, 409/12, 409/14, 487/04		A1	(11) International Publication Number: WO 00/31032 (43) International Publication Date: 2 June 2000 (02.06.00)
(21) International Application Number: PCT/EP99/08665 (22) International Filing Date: 11 November 1999 (11.11.99) (30) Priority Data: 60/109,297 20 November 1998 (20.11.98) US (71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH]; 124, Grenzacherstrasse, CH-4070 Basle (CH). (72) Inventors: ROGERS, Daniel, Harry; 8567 Hydra Lane, San Diego, CA 92126 (US). SAUNDERS, John; 5440 Caminico Exquisito, San Diego, CA 92130 (US). WILLIAMS, John, Patrick; 12066 Cassini Court, San Diego, CA 92131 (US). (74) Agent: BRAUN, Axel; 124 Grenzacherstrasse, CH-4070 Basle (CH).			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.

(54) Title: PYRROLIDINE DERIVATIVES-CCR-3 RECEPTOR ANTAGONISTS



(57) Abstract

This invention relates to certain 3-aminomethylpyrrolidine derivatives of Formula (I), wherein: Z is -N- or -(N⁺)⁻ wherein R is alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxyalkyl, alkoxyalkyl, or cyanoalkyl, and X⁻ is a pharmaceutically acceptable counterion; Ar¹ and Ar² are, independently of each other, aryl or heteroaryl; Q is straight or branched alkylene with 1-3 carbon atoms; R¹ is hydrogen or alkyl; A is either: (I) -N(R²)C(O)- when: B is: (i) an alkylene with 1-4 carbon atoms inclusive wherein one of the carbon atoms may optionally be replaced by a group selected from -C(O)-, -N(R⁴)-, -O-, -S(O)_n- (where n is 0, 1 or 2), -NR⁵C(O)- and -N(R⁶)SO₂-; or (ii) an alkynylene chain; or (II) a group selected from -N(R²)C(S)-, -N(R²)C(O)N(R³)-, -N(R²)C(S)N(R³)-, -N(R²)SO₂-N(R³)-, -N(R²)C(O)O-, and -OC(O)N(R³)- when: B is: (i) a bond; (ii) an alkylene chain of 1-4 carbon atoms inclusive wherein one of the carbon atoms may optionally be replaced by a group selected from -C(O)-, -N(R⁴)-, -O-, -S(O)_n- (where n is 0, 1 or 2), -NR⁵C(O)- and -N(R⁶)SO₂-; (iii) an alkenylene chain; or (iv) an alkynylene chain; that are CCR-3 receptor antagonists, pharmaceutical compositions containing them, methods for their use and methods for preparing these compounds.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

Pyrrolidine derivatives- CCR-3 Receptor Antagonists

This invention relates to certain pyrrolidine derivatives that are CCR-3
receptor antagonists, pharmaceutical compositions containing them, methods for their
5 use and methods for preparing these compounds.

Tissue eosinophilia is a feature of a number of pathological conditions such as
asthma, rhinitis, eczema and parasitic infections ((see Bousquet, J. et al. *N. Eng. J.*
Med. 323: 1033-1039 (1990) and Kay, A.B. and Corrigan, C.J. *Br. Med. Bull.* 48:51-
10 64 (1992)). In asthma, eosinophil accumulation and activation are associated with
damage to bronchial epithelium and hyperresponsiveness to constrictor mediators.
Chemokines such as RANTES, eotaxin and MCP-3 are known to activate eosinophils
((see Baggiolini, M. and Dahinden, C.A. *Immunol. Today.* 15:127-133 (1994), Rot, A.
M. et al. *J. Exp. Med.* 176, 1489-1495 (1992) and Ponath, P.D. et al. *J. Clin. Invest.*,
15 Vol. 97, #3, 604-612 (1996)). However, unlike RANTES and MCP-3 which also
induce the migration of other leukocyte cell types, eotaxin is selectively chemotactic
for eosinophils ((see Griffith-Johnson, D.A et al. *Biochem. Biophys. Res. Commun.*
197:1167 (1993) and Jose, P.J. et al. *Biochem. Biophys. Res. Commun.* 207, 788
(1994)). Specific eosinophil accumulation was observed at the site of administration
20 of eotaxin whether by intradermal or intraperitoneal injection or aerosol inhalation
((see Griffith-Johnson, D.A et al. *Biochem. Biophys. Res. Commun.* 197:1167 (1993);
Jose, P.J. et al. *J. Exp. Med.* 179, 881-887 (1994); Rothenberg, M.E. et al. *J. Exp.*
Med. 181, 1211 (1995) and Ponath, P.D. *J. Clin. Invest.*, Vol. 97, #3, 604-612
(1996)).

25 Glucocorticoids such as dexamethasone, methprednisolone and
hydrocortisone have been used for treating many eosinophil-related disorders,


including bronchial asthma ((R. P. Schleimer et. al., *Am. Rev. Respir. Dis.*, 141, 559 (1990)). The glucocorticoids are believed to inhibit IL-5, IL-3 mediated eosinophil survival in these diseases. However, prolonged use of glucocorticoids can lead to side effects such as glaucoma, osteoporosis and growth retardation in the patients ((see Hanania N.A et al., *J. Allergy and Clin. Immunol.*, Vol. 96, 571-579 (1995) and Saha M. T. et al, *Acta Paediatrica*, Vol. 86, #2, 138-142 (1997)). It is therefore desirable to have an alternative means of treating eosinophil related diseases without incurring these undesirable side effects.

Recently, the CCR-3 receptor was identified as a major chemokine receptor that eosinophils use for their response to eotaxin, RANTES and MCP-3. When transfected into a murine pre- β lymphoma line, CCR-3 bound eotaxin, RANTES and MCP-3 and conferred chemotactic responses on these cells to eotaxin, RANTES and MCP-3 ((see Ponath. P.D. et al. *J. Exp. Med.* 183, 2437-2448 (1996)). The CCR-3 receptor is expressed on the surface of eosinophils, T-cells (subtype Th-2), basophils and mast cells and is highly selective for eotaxin. Studies have shown that pretreatment of eosinophils with an anti-CCR-3 mAb completely inhibits eosinophil chemotaxis to eotaxin, RANTES and MCP-3 ((see Heath H. et al. *J. Clin. Invest.*, Vol. 99, #2, 178-184 (1997)).

Blocking the ability of the CCR-3 receptor to bind RANTES, MCP-3 and eotaxin and thereby preventing the recruitment of eosinophils should provide for the treatment of eosinophil-mediated inflammatory diseases:

The present invention therefore concerns novel pyrrolidine derivatives which are capable of inhibiting the binding of eotaxin to the CCR-3 receptor and thereby provide a means of combating eosinophil induced diseases, such as asthma.

In a first aspect, this invention provides compounds selected from the group of compounds represented by Formula (I):



5 wherein:

Z is -N- or -(N⁺R)- X⁻ wherein R is alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxyalkyl, alkoxycarbonylalkyl, or cyanoalkyl, and X⁻ is a pharmaceutically acceptable counterion;

Ar¹ and Ar² are, independently of each other, aryl or heteroaryl;

10

Q is straight or branched alkylene with 1-3 carbon atoms;

R¹ is hydrogen or alkyl;

A is either:

(I) -N(R²)C(O)- when:

B is:

15

(i) an alkylene with 1-4 carbon atoms inclusive wherein one of the carbon atoms may optionally be replaced by a group selected from -C(O)-, -N(R⁴)-, -O-, -S(O)_n- (where n is 0, 1 or 2), -NR⁵C(O)- and -N(R⁶)SO₂-; or

20

(ii) an alkynylene chain;

wherein:

R² is hydrogen, alkyl, acyl, haloalkyl, heteroalkyl, heterocyclalkyl, or -(alkylene)-C(O)-Z' where Z' is alkyl,

haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, or mono- or disubstituted amino; and

R^4 , R^5 and R^6 are, independently of each other, hydrogen, alkyl, acyl, haloalkyl, heteroalkyl, heterocyclalkyl, or -(alkylene)-C(O)-Z' where Z' is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, or mono- or disubstituted amino; or

(II) a group selected from -N(R^2)C(S)-, -N(R^2)C(O)N(R^3)-, -N(R^2)C(S)N(R^3)-, -N(R^2)SO₂-, -N(R^2)SO₂N(R^3)-, -N(R^2)C(O)O-, and -OC(O)N(R^3)- when:

B is:

- (i) a bond;
- (ii) an alkylene chain of 1-4 carbon atoms inclusive wherein one of the carbon atoms may optionally be replaced by a group selected from -C(O)-, -N(R^4)-, -O-, -S(O)_n- (where n is 0, 1 or 2), -NR⁵C(O)- and -N(R^6)SO₂-;
- (iii) an alkenylene chain; or
- (iv) an alkynylene chain;

wherein:

R^3 is hydrogen, alkyl, acyl, haloalkyl, heteroalkyl, heterocyclalkyl, or -(alkylene)-C(O)-Z' where Z' is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, or mono- or disubstituted amino; and

R^2 , R^4 , R^5 and R^6 are as defined above; and

prodrugs, individual isomers, mixtures of isomers and pharmaceutically acceptable salts thereof.

In a second aspect, this invention provides pharmaceutical compositions containing a therapeutically effective amount of a compound of Formula (I) or its pharmaceutically acceptable salt and a pharmaceutically acceptable excipient.

5 In a third aspect, this invention provides a method of treatment of a disease in a mammal treatable by administration of a CCR-3 receptor antagonist, comprising administration of a therapeutically effective amount of a compound of Formula (I) or its pharmaceutically acceptable salt. The disease states include respiratory diseases such as asthma.

10 In a fourth aspect, this invention provides a process for preparing compounds of Formula (I).

15 Unless otherwise stated, the following terms used in the specification and claims have the meanings given below:

"Alkyl" means a linear saturated monovalent hydrocarbon radical of one to six carbon atoms or a branched saturated monovalent hydrocarbon radical of three to six carbon atoms, e.g., methyl, ethyl, propyl, 2-propyl, pentyl, and the like.

20 "Alkenyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, e.g., ethenyl, propenyl, and the like.

25 "Alkylene" means a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms, e.g., methylene, ethylene, propylene, 2-methylpropylene, pentylene, and the like.

"Alkenylene" means a linear divalent hydrocarbon radical of two to six carbon atoms or a branched divalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, e.g., ethenylene, 2,4-pentadienylene, and the like.

5 "Alkynylene" means a linear divalent hydrocarbon radical of two to six carbon atoms or a branched divalent hydrocarbon radical of three to six carbon atoms, containing at least one triple bond, e.g., ethynylene, propynylene, and the like.

10 "Acyl" means a radical $-C(O)R$ where R is alkyl, haloalkyl, optionally substituted phenyl, optionally substituted phenylalkyl, optionally substituted heteroaralkyl or optionally substituted heteroaryl, e.g., acetyl, benzoyl, thenoyl, and the like.

15 "Halo" means fluoro, chloro, bromo or iodo, preferably fluoro and chloro.

"Haloalkyl" means alkyl substituted with one or more same or different halo atoms, e.g., $-CH_2Cl$, $-CF_3$, $-CH_2CF_3$, $-CH_2CCl_3$, and the like.

20 "Cycloalkyl" means a saturated monovalent cyclic hydrocarbon radical of three to six ring carbons, e.g., cyclopropyl, cyclohexyl, and the like.

"Monosubstituted-amino" means a radical $-NHR$ where R is alkyl, heteroalkyl, haloalkyl, optionally substituted phenyl or optionally substituted heteroaryl, e.g., methylamino, (1-methylethyl)amino, phenylamino, and the like.

25 "Disubstituted-amino" means a radical $-NRR'$ where R and R' are independently alkyl, heteroalkyl, haloalkyl, optionally substituted phenyl, or optionally substituted heteroaryl. Representative examples include, but are not limited to, dimethylamino, methylethylamino, di(1-methylethyl)amino, and the like.

"Aryl" means a monovalent monocyclic or bicyclic aromatic hydrocarbon radical of 6 to 10 ring atoms. The aryl ring may be optionally substituted independently with one or more substituents, preferably one, two or three substituents selected from alkyl, haloalkyl, alkylthio, heteroalkyl, alkoxy, cycloalkyl, cycloalkylalkyl, alkenyl, halo, cyano, nitro, optionally substituted phenyl, optionally substituted phenylalkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, amino, monosubstituted amino, disubstituted amino, hydroxylamino, -OR [where R is hydrogen, haloalkyl, optionally substituted phenyl, optionally substituted phenylalkyl, optionally substituted heteroaryl or optionally substituted heteroaralkyl], $-S(O)_nR$ [where n is an integer from 0 to 2 and R is alkyl, haloalkyl, optionally substituted phenyl, optionally substituted phenylalkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, amino, mono- or disubstituted amino], $-NRC(O)R'$ (where R is hydrogen or alkyl and R' is hydrogen, alkyl, heteroalkyl, haloalkyl, optionally substituted phenyl, or mono- or disubstituted amino), $-NRSO_2R'$ (where R is hydrogen or alkyl and R' is alkyl, amino, mono- or disubstituted amino), $-C(O)R$ (where R is hydrogen, alkyl, haloalkyl, optionally substituted phenyl or optionally substituted heteroaryl), $-COOR$ (where R is hydrogen, alkyl, optionally substituted phenyl, optionally substituted phenylalkyl, optionally substituted heteroaryl or optionally substituted heteroaralkyl), $-(alkylene)-COOR$ (where R is hydrogen, alkyl, optionally substituted phenyl, optionally substituted phenylalkyl, optionally substituted heteroaryl or optionally substituted heteroaralkyl), $alkylenedioxy$, $oxy-C2-C3-alkylene$, $-CONR'R''$ or $-(alkylene)-CONR'R''$ (where R' and R'' are independently selected from hydrogen, alkyl, haloalkyl, optionally substituted phenyl, optionally substituted phenylalkyl, optionally substituted heteroaryl or optionally substituted heteroaralkyl). More specifically the term aryl includes, but is not limited to, phenyl, 1-naphthyl, 2-naphthyl, and derivatives thereof.

"Alkylenedioxy" is a divalent substituent of the formula $[-O-(CH_2)_n-O-]$ (where n is 1 or 2) which is attached to two adjacent carbon atoms of phenyl, e.g., methylenedioxy-phenyl or ethylenedioxyphenyl.

5 "Oxy-C2-C3-alkylene" means a divalent substituent of the formula $[-O-(CH_2)_n-]$ (where n is 2 or 3) which is attached to two adjacent carbon atoms of phenyl, e.g., oxyethylene or oxypropylene. An example of oxy-C2-C3-alkylenephenyl is 2,3-dihydrobenzofuranyl.

10 "Optionally substituted phenyl" means a phenyl group which is optionally substituted independently with one, two or three substituents selected from alkyl, haloalkyl, halo, nitro, cyano, -OR (where R is hydrogen or alkyl), -NRR' (where R and R' are independently of each other hydrogen or alkyl), -COOR (where R is hydrogen or alkyl) or -CONR'R'' (where R' and R'' are independently selected
15 from hydrogen or alkyl).

"Heteroaryl" means a monovalent monocyclic or bicyclic aromatic radical of 5 to 10 ring atoms containing one, two, or three ring heteroatoms selected from N, O, or S, the remaining ring atoms being C. The aromatic radical is optionally fused to a
20 phenyl or an optionally substituted heteroaryl ring or it is optionally substituted independently with one or more substituents, preferably one or two substituents selected from alkyl, haloalkyl, heteroalkyl, alkoxy, halo, cyano, nitro, aryl, optionally substituted heteroaryl, amino, monosubstituted amino, disubstituted amino, hydroxyamino, -OR [where R is hydrogen, haloalkyl, or optionally substituted
25 phenyl], $-S(O)_nR$ [where n is an integer from 0 to 2 and R is alkyl, haloalkyl, optionally substituted phenyl, amino, mono or disubstituted amino], $-C(O)R$ (where R is hydrogen, alkyl, haloalkyl or optionally substituted phenyl), -COOR (where R is hydrogen, alkyl or optionally substituted phenyl), $-(alkylene)-COOR$ (where R is hydrogen, alkyl or optionally substituted phenyl), methylenedioxy, 1,2-ethylenedioxy,
30 $-CONR'R''$ or $-(alkylene)-CONR'R''$ (where R' and R'' are independently selected

from hydrogen, alkyl, haloalkyl, or optionally substituted phenyl). More specifically the term heteroaryl includes, but is not limited to pyridyl, pyrrolyl, thiophene, pyrazolyl, thiazolyl, imidazolyl, pyrimidinyl, thiadiazolyl, indolyl, carbazolyl, azaindolyl, benzofuranyl, benzimidazolyl, benzthiazolyl, quinoxaliny, benzotriazolyl, benzisoxazolyl, purinyl, quinolinyl, isoquinolinyl, benzopyranyl, 9H-1,3,4,9-tetraazafluorene, and derivatives thereof.

"Optionally substituted heteroaryl" means a pyridyl, pyrrolyl, thiophene, pyrazolyl, thiazolyl, imidazolyl, pyrimidinyl, thiadiazolyl, indolyl, carbazolyl, azaindolyl, benzofuranyl, benzimidazolyl, benzthiazolyl, quinoxaliny, benzotriazolyl, benzisoxazolyl, purinyl, quinolinyl, isoquinolinyl, benzopyranyl, 9H-1,3,4,9-tetraazafluorene ring which is optionally substituted independently with one, two or three substituents selected from alkyl, haloalkyl, halo, nitro, cyano, -OR (where R is hydrogen or alkyl), -NRR' (where R and R' are independently of each other hydrogen or alkyl), -COOR (where R is hydrogen or alkyl) or -CONR'R" (where R' and R" are independently selected from hydrogen or alkyl).

"Heterocycle" or "Heterocyclyl" means a saturated or unsaturated cyclic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatoms selected from N, O, or S(O)_n (where n is an integer from 0 to 2), the remaining ring atoms being C, where one or two C atoms may optionally be replaced by a carbonyl group. The heterocycle ring may be optionally substituted independently with one, two or three substituents selected from alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, halo, cyano, acyl, acylamino, amino, monosubstituted amino, disubstituted amino, -COOR (where R is hydrogen or alkyl), -XR (where X is O or S(O)_n, where n is an integer from 0 to 2 and R is hydrogen, alkyl, haloalkyl, cycloalkyl, aralkyl, aryl, heteroaryl or heteroaralkyl), or -CONR'R" (where R' and R" are independently selected from hydrogen or alkyl).

Representative examples include, but are not limited to, tetrahydropyranyl, piperidino, piperazino, pyrrolidino, and the like.

"Heteroalkyl" means an alkyl, cycloalkyl, or cycloalkylalkyl radical as defined above, carrying a substituent containing a heteroatom selected from N, O, S(O)_n 5 where n is an integer from 0 to 2. Representative substituents include -NR^aR^b, -OR^a or -S(O)_nR^c, wherein n is an integer from 0 to 2, R^a is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, optionally substituted phenyl, optionally substituted phenylalkyl, optionally substituted heteroaryl, or -COR (where R is alkyl), R^b is 10 hydrogen, alkyl, -SO₂R (where R is alkyl or hydroxyalkyl), -SO₂NRR' (where R and R' are independently of each other hydrogen or alkyl), -CONR'R" (where R' and R" are independently selected from hydrogen or alkyl) and R^c is hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, optionally substituted phenyl, optionally substituted heteroaryl, amino, monosubstituted amino, or disubstituted amino. Representative 15 examples include, but are not limited to 2-methoxyethyl, 2-hydroxyethyl, 2-aminoethyl, 2-dimethylaminoethyl, benzyloxymethyl, thiophen-2-ylthiomethyl, and the like.

"Hydroxyalkyl" means a linear monovalent hydrocarbon radical of two to six 20 carbon atoms or a branched monovalent hydrocarbon radical of three or six carbons substituted with one or two hydroxy groups, provided that if two hydroxy groups are present they are not both on the same carbon atom. Representative examples include, but are not limited to, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 1-(hydroxymethyl)-2-methylpropyl, 2-hydroxybutyl, 3-hydroxybutyl, 4-hydroxybutyl, 25 2,3-dihydroxypropyl, 1-(hydroxymethyl)-2-hydroxyethyl, 2,3-dihydroxybutyl, 3,4-dihydroxybutyl and 2-(hydroxymethyl)-3-hydroxypropyl, preferably 2-hydroxyethyl, 2,3-dihydroxypropyl, and 1-(hydroxymethyl)-2-hydroxyethyl.

"Cycloalkylalkyl" means a radical $-R^aR^b$ where R^a is an alkylene group and R^b is a cycloalkyl group as defined above e.g., cyclopropylmethyl, cyclohexylpropyl, 3-cyclohexyl-2-methylpropyl, and the like.

5 "Aralkyl" means a radical $-R^aR^b$ where R^a is an alkylene group and R^b is an aryl group as defined above e.g., benzyl, phenylethyl, 3-(3-chlorophenyl)-2-methylpentyl, and the like.

10 "Alkoxy" or "haloalkyloxy" means a radical $-OR$ where R is an alkyl or haloalkyl, respectively as defined above e.g., methoxy, and the like.

"Optional" or "optionally" means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For
15 example, "heterocyclo group optionally mono- or di- substituted with an alkyl group" means that the alkyl may but need not be present, and the description includes situations where the heterocyclo group is mono- or disubstituted with an alkyl group and situations where the heterocyclo group is not substituted with the alkyl group.

20 "Amino-protecting group" refers to those organic groups intended to protect nitrogen atoms against undesirable reactions during synthetic procedures e.g., benzyl, benzyloxycarbonyl (CBZ), t-butoxycarbonyl (BOC), trifluoroacetyl, and the like.

25 Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers". Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers" and those that are non-superimposable mirror images of each other are termed "enantiomers". When a compound has an asymmetric center, for
30 example, if a carbon atom is bonded to four different groups, a pair of enantiomers is

possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the *R*- and *S*-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture".

The compounds of this invention may possess one or more asymmetric centers; such compounds can therefore be produced as individual (*R*)- or (*S*)-stereoisomers or as mixtures thereof. For example, if the R^1 substituent in a compound of Formula (I) is alkyl, then the carbon to which it is attached is an asymmetric center and the compound of Formula (I) can exist as an (*R*)- or (*S*)-stereoisomer. Unless indicated otherwise, the description or naming of a particular compound in the specification and claims is intended to include both individual enantiomers and mixtures, racemic or otherwise, thereof. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art (see discussion in Chapter 4 of "Advanced Organic Chemistry", 4th edition J. March, John Wiley and Sons, New York, 1992).

A "pharmaceutically acceptable excipient" means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes an excipient that is acceptable for veterinary use as well as human pharmaceutical use. "A pharmaceutically acceptable excipient" as used in the specification and claims includes both one and more than one such excipient.

A "pharmaceutically acceptable counterion" means an ion having a charge opposite to that of the substance with which it is associated and that is pharmaceutically acceptable. Representative examples include, but are not limited to,

chloride, bromide, iodide, methanesulfonate, p-tolylsulfonate, trifluoroacetate, acetate, and the like.

A "pharmaceutically acceptable salt" of a compound means a salt that is
5 pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. Such salts include:

(1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed
10 with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid,
15 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid,
20 glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or

(2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion; an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine,
25 diethanolamine, triethanolamine, tromethamine, *N*-methylglucamine, and the like.

"Leaving group" has the meaning conventionally associated with it in synthetic organic chemistry i.e., an atom or group capable of being displaced by a nucleophile and includes halogen, alkanesulfonyloxy, arenesulfonyloxy, ester, or

amino such as chloro, bromo, iodo, mesyloxy, tosyloxy, trifluorosulfonyloxy, methoxy, N,O-dimethylhydroxylamino, and the like.

"Pro-drugs" means any compound which releases an active parent drug according to Formula (I) *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of Formula (I) are prepared by modifying functional groups present in the compound of Formula (I) in such a way that the modifications may be cleaved *in vivo* to release the parent compound. Prodrugs include compounds of Formula (I) wherein a hydroxy, sulfhydryl or amino group in compound (I) is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino, or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to esters (e.g., acetate, formate, and benzoate derivatives), carbamates (e.g., *N,N*-dimethylaminocarbonyl) of hydroxy functional groups in compounds of Formula (I), and the like.

15

"Treating" or "treatment" of a disease includes:

- (1) preventing the disease, i.e. causing the clinical symptoms of the disease not to develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease,
- (2) inhibiting the disease, i.e., arresting or reducing the development of the disease or its clinical symptoms, or
- (3) relieving the disease, i.e., causing regression of the disease or its clinical symptoms.


25

A "therapeutically effective amount" means the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

30

The nomenclature used in this application is generally based on the IUPAC recommendations.

The pyrrolidine ring is numbered as follows:



and the compounds of the invention are named as:

A compound of Formula (I) where Z is -N-, Ar¹ is 3,4-dichlorophenyl, Q is -CH₂-, R¹ is hydrogen, A is -NHCO-, B is -(CH₂)₂- and Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl, is named *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-ylmethyl]-3-[5-(4-methoxyphenyl)-pyrimidin-2-yl]propionamide.

A compound of Formula (I) where Z is -N-, Ar¹ is 3,4-dichlorophenyl, Q is -CH₂-, R¹ is hydrogen, A is -NHCO-, B is -CH₂S- and Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl, is named *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-[5-(4-methoxyphenyl)-pyrimidin-2-ylsulfanyl]acetamide.

A compound of Formula (I) where Z is -N-, Ar¹ is 3,4-dichlorophenyl, Q is -CH₂-, R¹ is hydrogen, A is -NHCO-, B is -CH₂O- and Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl, is named *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-[5-(4-methoxyphenyl)-pyrimidin-2-yloxy]acetamide.

A compound of Formula (I) where Z is -N-, Ar¹ is 3,4-dichlorophenyl, Q is -CH₂-, R¹ is hydrogen, A is -NHCO-, B is -CH₂N- and Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl, is named *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-[5-(4-methoxyphenyl)-pyrimidin-2-ylamino]acetamide.


A compound of Formula (I) where Z is -N-, Ar¹ is 3,4-methylenedioxyphenyl, Q is -CH₂-, R¹ is hydrogen, A is -NHCO-, B is -CH₂S- and Ar² is 5-(4-methoxyphenyl)-pyrimidin-2-yl, is named *N*-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-ylsulfanyl]acetamide.

A compound of Formula (I) where Z is -N-, Ar¹ is 3,4-dichlorophenyl, Q is -CH₂-, R¹ is hydrogen, A is -NHCO-, B is -CH₂- and Ar² is 4-methoxyphenyl, is named *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-(4-methoxyphenyl)acetamide.

A compound of Formula (I) where Z is -N-, Ar¹ is 3,4-dichlorophenyl, Q is -CH₂-, R¹ is hydrogen, A is -NHCONH-, B is a bond and Ar² is 4-methoxyphenyl, is named 1-[1-(3,4-dichlorobenzyl)pyrrolidin-3-ylmethyl]-3-(4-methoxyphenyl)urea.

Representative compounds of this invention are as follows:

- I. Representative compounds of Formula (I) where Z is -N-, R¹ is hydrogen and A is -NHCO-



CPD #	Ar ¹	Q	B	Ar ²	Mass. Spec. m/e
1.	2,3-dichlorophenyl	CH ₂	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl	519
2.	3,4-methylenedioxy-phenyl	CH ₂	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl . HCl	493
3.	2,3-dichlorophenyl	CH ₂	(CH ₂) ₂	5-(4-methoxyphenyl)pyrimidin-2-yl	500
4.	3,4-dichlorophenyl	CH ₂	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl	519
5.	2,3-dichlorophenyl	CH ₂	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl . HCl	
6.	2,3-dichlorophenyl	CH ₂	(CH ₂) ₂	5-(4-dimethylaminophenyl)pyrimidin-2-yl	514
7.	2,3-dichlorophenyl	CH ₂	(CH ₂) ₂	5-(4-methoxyphenyl)pyrimidin-2-yl	499
8.	3,4-methylenedioxy-phenyl	CH ₂	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl	494
9.	2,3-dichlorophenyl	CH ₂	CH ₂ O	5-(4-methoxyphenyl)pyrimidin-2-yl	502

CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
10.	3,4-methylenedioxy-phenyl	CH ₂	(CH ₂) ₂	5-(4-methoxyphenyl)pyrimidin-2-yl	476
11.	3-chlorophenyl	CH ₂	(CH ₂) ₂	5-(4-methoxyphenyl)pyrimidin-2-yl	466
12.	3,4-dichlorophenyl	CH ₂	CH ₂ S	9H-1,3,4,9-tetraaza-fluoren-2-yl .TFA	501
13.	3,4-methylenedioxy-phenyl	CH ₂	CH ₂ S	5-phenylpyrimidin-2-yl	464
14.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂	5-(4-methoxyphenyl)pyrimidin-2-yl	500
15.	3,4-dichlorophenyl	CH ₂	CH ₂ NH ₂ SO ₂	phenyl .TFA	
16.	2,3-dichlorophenyl	CH ₂	CH ₂ O	5-(4-methylthiophenyl)pyrimidin-2-yl	518
17.	3-chlorophenyl	CH ₂	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl	484
18.	3,4-dichlorophenyl	CH ₂	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl	519
19.	2,3-dichlorophenyl	CH ₂	CH ₂ O	5-benzofuran-2-ylpyrimidin-2-yl	512
20.	2,3-dichlorophenyl	CH ₂	CH ₂ S	5-phenylpyrimidin-2-yl	489
21.	3,4-methylenedioxy-phenyl	CH ₂	CH ₂ O	5-(4-methoxyphenyl)pyrimidin-2-yl	477
22.	2,3-methylenedioxy-phenyl	CH ₂	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl	494
23.	3,4-dichlorophenyl	CH ₂	CH ₂ O	5-(4-methoxyphenyl)pyrimidin-2-yl	502
24.	3-chlorophenyl	CH ₂	(CH ₂) ₂	5-(4-dimethylaminophenyl)pyrimidin-2-yl	479

CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
25.	2,3-dichlorophenyl	CH ₂	CH ₂ S	5-(4-methylphenyl)pyrimidin-2-yl	503
26.	3,4-dichlorophenyl	CH ₂	CH ₂ S	5-phenylpyrimidin-2-yl	
27.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂	5-(4-dimethylaminophenyl)pyrimidin-2-yl	514
28.	2,3-dichlorophenyl	CH ₂	CH ₂ O	5-(4-methylphenyl)pyrimidin-2-yl	486
29.	3,4-methylenedioxy-phenyl	CH ₂	CH ₂ S	5-phenylpyrimidin-2-yl	464
30.	2,3-dichlorophenyl	CH ₂	CH ₂ O	5-(4-fluorophenyl)pyrimidin-2-yl	490
31.	3-chlorophenyl	CH ₂	CH ₂ S	5-phenylpyrimidin-2-yl	454
32.	2,3-dichlorophenyl	CH ₂	CH ₂ S	5-phenylpyrimidin-2-yl	489
33.	3,4-methylenedioxy-phenyl	CH ₂	CH ₂ O	5-(4-methoxyphenyl)pyrimidin-2-yl	477
34.	3,4-methylenedioxy-phenyl	CH ₂	CH ₂ S	5-(4-chlorophenyl)pyrimidin-2-yl	498
35.	naphth-2-yl	CH ₂	(CH ₂) ₂	5-(4-methoxyphenyl)pyrimidin-2-yl	482
36.	2,3-dichlorophenyl	CH ₂	CH ₂ O	5-(4-vinylphenyl)pyrimidin-2-yl	498
37.	2,3-dichlorophenyl	CH ₂	CH ₂ S	5-(4-chlorophenyl)pyrimidin-2-yl	523
38.	3,4-methylenedioxy-phenyl	CH ₂	CH ₂ S	5-(4-methylphenyl)pyrimidin-2-yl	478
39.	2,3-dichlorophenyl	CH ₂	(CH ₂) ₂	5-(4-methylphenyl)pyrimidin-2-yl	484

CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
40.	2,3-dichlorophenyl	CH ₂	CH ₂ N	5-(4-methoxyphenyl)pyrimidin-2-yl	
41.	2,3-dichlorophenyl	CH ₂	(CH ₂) ₂	1-(4-aminosulfonylphenyl)pyrazol-4-yl	537
42.	3,4-dichlorophenyl	CH ₂	CH ₂ S	pyrimidin-2-yl . HCl	
43.	3,4-dichlorophenyl	CH ₂	CH ₂ S	pyridin-4-yl . TFA	410
44.	3,4-dichlorophenyl	CH ₂	CH ₂ S	4-amino-5-(3-chlorophenyl)pyrimidin-2-yl	536
45.	3,4-dichlorophenyl	CH ₂	CH ₂ S	quinoxalin-2-yl	461
46.	3,4-methylenedioxy-phenyl	CH ₂	CH ₂ O	5-(4-methylthiophenyl)pyrimidin-2-yl	493
47.	3,4-dichlorophenyl	CH ₂	CH ₂ S	naphth-2-yl . TFA	459
48.	3,4-dichlorophenyl	CH ₂	CH ₂	1-phenyl-5-methylpyrazol-4-yl . TFA	457
49.	3,4-methylenedioxy-phenyl	CH ₂	(CH ₂) ₂	5-(2-methylphenyl)pyrimidin-2-yl	460
50.	naphth-2-yl	CH ₂	(CH ₂) ₂	5-(2-methylphenyl)pyrimidin-2-yl	466
51.	2,3-dichlorophenyl	CH ₂	(CH ₂) ₂	1-(4-methylphenyl)pyrazol-4-yl	537
52.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂	2-bromo-4,5-dimethoxyphenyl . TFA	383
53.	3,4-dichlorophenyl	CH ₂	CH ₂	thiophen-3-yl . TFA	
54.	3,4-methylenedioxy-	CH ₂	CH ₂ O	5-(benzofuran-2-yl)pyrimidin-2-yl	487


CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
	phenyl				
55.	3,4-dichlorophenyl	CH ₂	CH ₂ S	4-methylpyrimidin-2-yl . TFA	425
56.	3,4-dichlorophenyl	CH ₂	CH ₂ S	4-aminophenyl . TFA	
57.	2,3-dichlorophenyl	CH ₂	(CH ₂) ₂	1-[4-(2-propylphenyl)]pyrazol-4-yl	461
58.	6-chloro-3,4-methylenedioxyphenyl	CH ₂	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl	528
59.	3,4-dichlorophenyl	CH ₂	CH ₂ O	4-benzoyloxyphenyl . TFA	499
60.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₃ CO	thiophen-2-yl . TFA	439
61.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂	1-benzylpyrazol-4-yl	452
62.	3,4-dichlorophenyl	CH ₂	CH ₂ O	4-acetylphenyl . TFA	435
63.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂	3-hydroxyphenyl . TFA	407
64.	3,4-dichlorophenyl	CH ₂	CH ₂	3-(thiophen-2-yl)pyrazol-1-yl	449
65.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₃	3,4-dimethoxyphenyl . TFA	465
66.	3,4-dichlorophenyl	CH ₂	CH ₂	5,6-dimethylbenzimidazol-1-yl . TFA	445
67.	3,4-dichlorophenyl	CH ₂	CH ₂ S	5-trifluoromethylpyridin-2-yl	
68.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₃ O	4-chloro-3-methylphenyl . TFA	

CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
69.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂ CO	naphth-1-yl . TFA	469
70.	4-fluoro-3-methoxy-phenyl	CH ₂	CH ₂ O	5-(4-methoxyphenyl)pyrimidin-2-yl	481
71.	2,3-dichlorophenyl	CH ₂	(CH ₂) ₂	1-(2,6-dimethylpyrimidin-4-yl)pyrazol-4-yl	488
72.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₃ O	2,4-dichlorophenyl . TFA	490
73.	3,4-dichlorophenyl	CH ₂	CH ₂ S	4,6-dimethoxy-5-phenylpyrimidin-2-yl	519
74.	3,4-dichlorophenyl	(CH ₂) ₃	CH ₂ S	5-(4-methoxyphenyl)pyrimidin-2-yl	546
75.	3,4-dichlorophenyl	CH ₂	CH ₂	indol-1-yl . TFA	416
76.	3,4-dichlorophenyl	CH ₂	CH ₂ S	2-(pyridin-2-yl)-4-trifluoromethylpyrimidin-6-yl . TFA	556
77.	3,4-dichlorophenyl	CH ₂	CH ₂ N(CH ₃)CO	phenyl	
78.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂ CO	4-methoxynaphth-1-yl . TFA	
79.	3,4-dichlorophenyl	CH ₂	CH ₂ O	4-hydroxymethylphenyl . TFA	423
80.	3,4-ethylenedioxy-phenyl	CH ₂	CH ₂ S	5-(4-methylphenyl)pyrimidin-2-yl	492
81.	3,4-dichlorophenyl	CH ₂	CH ₂ SO ₂	4-methylphenyl . TFA	455
82.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂	1-(quinoxalin-2-yl)pyrazol-4-yl	510
83.	3,4-dichlorophenyl	CH ₂	CH ₂ S	5-(4-chlorophenyl)pyrimidin-4-yl	

CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
84.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂ CO	4-methylsulfonylphenyl.TFA	497
85.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂ CO	3,6-dimethoxynaphth-2-yl.TFA	529
86.	3,4-dichlorophenyl	CH ₂	CH(CH ₃)O	1-acetylnaphth-2-yl.TFA	499
87.	2,3-dichlorophenyl	CH ₂	CH ₂	3-(2,5-difluoro-4-chlorophenylcarbonylamino)phenyl	
88.	2,3-dichlorophenyl	CH ₂	CH ₂	3-(2,3,4,5-tetrafluorophenylcarbonylamino)phenyl	
89.	2,3-dichlorophenyl	CH ₂	CH ₂	3-(acetylamino)phenyl	
90.	2,3-dichlorophenyl	CH ₂	CH ₂	3-fluoro-4-hydroxyphenyl	411
91.	3,4-methylenedioxyphenyl	CH ₂	CH ₂	3-(2,4-dimethoxyphenylcarbonylamino)phenyl	
92.	3,4-dichlorophenyl	CH ₂	CH ₂	4-methylsulfonylphenyl	
93.	3,4-dichlorophenyl	CH ₂	*CH(CH ₃) * = (R) stereochem.	6-methoxynaphth-2-yl	471
94.	3,4-dichlorophenyl	CH ₂	CH ₂	5-benzoylthiophen-2-yl.TFA	487
95.	3,4-dichlorophenyl	CH ₂	CH ₂	3-fluoro-4-hydroxyphenyl	411
96.	3,4-methylenedioxyphenyl	CH ₂	CH ₂ S	9 H-1,3,4,9-tetraazafluoren-2-yl.TFA	477
97.	2,3-dichlorophenyl	CH ₂	CH ₂ O	5-(3,4-methylenedioxyphenyl)pyrimidin-2-yl	531

CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
98.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂	5-(3,4-methylenedioxyphenyl)pyrimidin-2-yl	529
99.	2,3-dichlorophenyl	CH ₂	(CH ₂) ₂	5-(3,4-methylenedioxyphenyl)pyrimidin-2-yl	529
100.	2,3-dichloro-thiophen-5-yl	CH ₂	CH ₂ S	5-(3,4-methylenedioxyphenyl)pyrimidin-2-yl	553
101.	2,3-dichlorophenyl	CH(CH ₃)	CH ₂ S	5-(3,4-methylenedioxy)pyrimidin-2-yl	561
102.	3,4-dichlorophenyl	CH ₂	CH ₂ S	5-(3,4-dimethoxyphenyl)pyrimidin-2-yl	547


II. Representative compounds of Formula (I) where Z is -N-, R¹ is hydrogen and A is -NHCONH-.



CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
103.	3,4-methylenedioxyphenyl	CH ₂	CH ₂	3-methylphenyl .TFA	382
104.	3,4-methylenedioxyphenyl	CH ₂	bond	3-methylphenyl .TFA	368
105.	3,4-dichlorophenyl	CH ₂	CH ₂	4-pyridyl	
106.	2,3-dichlorophenyl	CH ₂	CH ₂	4-methoxyphenyl	423
107.	3-biphenyl	CH ₂	bond	3-methylphenyl .TFA	400
108.	3,4-dichlorophenyl	CH ₂	CH ₂	2-pyridyl	
109.	3,4-dichlorophenyl	CH ₂	CH ₂	3-fluoro-5-trifluoromethylphenyl	
110.	3,4-dichlorophenyl	CH ₂	bond	2-methoxyphenyl	408
111.	3,4-dichlorophenyl	CH ₂	(CH ₂) ₂	phenyl .TFA	406


CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. - m/e
112.	3,4-dichlorophenyl	(CH ₂) ₂	bond	3-methoxyphenyl .TFA	422
113.	naphth-2-yl	CH ₂	CH ₂	4-methoxyphenyl	405
114.	3,4-dichlorophenyl	CH ₂	bond	3-methoxycarbonylphenyl	436
115.	3,4-dichlorophenyl	(CH ₂) ₃	bond	3-chlorophenyl .TFA	422
116.	3,4-dichlorophenyl	(CH ₂) ₂	(CH ₂) ₂	phenyl .TFA	420
117.	3,4-dichlorophenyl	CH ₂	bond	4-phenoxyphenyl	484
118	5-bromothiophen-2-yl	CH ₂	bond	3-methoxyphenyl	440
119	3,4-dichlorophenyl	CH ₂	-CO-	phenyl .TFA	

III. Representative compounds of Formula (I) where Z is -N-, R¹ is hydrogen and A is -NHC(S)NH-.




CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
120.	3,4-dichlorophenyl	CH ₂	bond	4-dimethylaminophenyl	
121.	3-phenyl	(CH ₂) ₃	bond	4-dimethylaminophenyl	400

IV. Representative compounds of Formula (I) where Z is -N-, R¹ is hydrogen and A is -NHSO₂-.



CPD #	Ar ¹	Q	B	Ar ²	Mass Spec. m/e
122.	3-(3,4-dichlorophenoxy)phenyl	CH ₂	CH ₂	phenyl	506
123.	8-carboxynaphth-1-yl	CH ₂	CH ₂	phenyl	482


V. Representative compounds of Formula (I) where Z is $-(NR^5)-$, X^- , R^1 is hydrogen and A is $-NHCONH-$.



CPD #	Ar ¹	Q	B	R	X ⁻	Ar ²
124.	3,4-dichlorophenyl	CH ₂	CH ₂ S	CH ₃	iodide	5-(4-methoxyphenyl)pyrimidin-2-yl

PREFERRED EMBODIMENTS

While the broadest definition of this invention is set forth above as the first aspect of the present invention, certain compounds of Formula (I) are preferred.



5

(I)

(I) One preferred group of compounds is that wherein:

R^1 is hydrogen; and

A is -NHCO-.

10

Within this group, a more preferred group of compounds is that wherein:

(a) Z is -N-; and

Ar^1 is a naphthyl or a substituted phenyl ring.

15

Within these preferred and more preferred groups, an even more preferred group of compounds is that wherein:

The stereochemistry at the C-3 carbon of the pyrrolidine ring is (S);

Q is a straight or branched alkylene chain of 1 to 3 carbon atoms, preferably

20

-CH₂-, -CH(CH₃)-, -C(CH₃)₂-;

B are -CH₂-;

Ar^1 is a phenyl ring substituted with one, two or three substituents selected from alkyl, cyano, nitro, halo, alkylendioxy, oxy-C2-C3-alkylene, alkoxy or phenoxy,

preferably a phenyl ring substituted with one or two substituents selected from methyl, chloro, fluoro, bromo, or methylenedioxy; and

Ar^2 is an aryl ring, preferably a phenyl ring which is optionally substituted with one, two, or three substituents selected from alkoxy, alkylthio, halo, amino, -

5 $NHC(O)R'$ (where R' is alkyl or optionally substituted phenyl), hydroxy, or $-SO_2Me$, preferably a phenyl ring optionally substituted with one, two, or three substituents selected from methyl, methylthio, hydroxy, methoxy, acetyl, chloro, fluoro, bromo, or -
 $NHC(O)R'$ (where R' is methyl or a phenyl ring optionally substituted with one, two or three substituents selected from methyl, methoxy, fluoro or chloro);

10

Within these preferred and more preferred groups, particularly preferred group of compounds is that wherein:

Ar^1 is 3-chlorophenyl, 4-chlorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl,
 15 3,4-ethylenedioxyphenyl, 2-naphthyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, 3,4-methylenedioxyphenyl, most preferably 2,3-dichlorophenyl, 3,4-dichlorophenyl or 3,4-methylenedioxyphenyl; and

Ar^2 is phenyl, 3-(2,5-difluoro-4-chlorophenylcarbonylamino)phenyl, 4-(2,3,4,5-tetrafluorophenylcarbonylamino)phenyl, 3-(2,4-dimethoxyphenylcarbonylamino)phenyl, 3-(phenylcarbonylamino)phenyl, 3-acetylaminophenyl, 3-fluoro-4-hydroxyphenyl, 3-fluorophenyl, 3-methoxyphenyl or 4-hydroxy-3-methoxyphenyl, most preferably 3-(2,5-difluoro-4-chlorophenylcarbonylamino)phenyl, 3-acetylaminophenyl, 3-(2,4-dimethoxyphenylcarbonylamino)phenyl or 3-(phenylcarbonylamino)phenyl.

25

(b) A second more preferred group of compounds in group (I) is that wherein:

Z is -N-; and

B is alkylene chain, preferably $-(CH_2)_2-$.

Within this more preferred group, an even more preferred group of compounds is that wherein:

The stereochemistry at the C-3 carbon of the pyrrolidine ring is (*S*);

Q is a straight or branched alkylene chain of 1 to 3 carbon atoms, preferably

5 -CH₂-, -CH(CH₃)-, -C(CH₃)₂-; and

Ar¹ is naphthyl or a substituted phenyl ring.

Within these more preferred and even more preferred groups, a particularly preferred group of compounds is that wherein:

10 Ar¹ is a phenyl ring substituted with one, two or three substituents selected from alkyl, cyano, nitro, halo, alkylenedioxy, oxy-C2-C3-alkylene, alkoxy or phenoxy, preferably one or two substituents selected from methyl, chloro, fluoro, bromo, methylenedioxy; more preferably 3-chlorophenyl, 4-chlorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 3,4-ethylenedioxyphenyl, 2-naphthyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, 3,4-methylenedioxyphenyl, most preferably 2,3-dichlorophenyl, 3,4-dichlorophenyl or 3,4-methylenedioxyphenyl.

20 Within the above preferred, more preferred group and particularly preferred groups, an even more preferred group of compounds is that wherein:

(i) Ar² is a pyrimidin-2-yl ring optionally substituted with alkyl or a phenyl ring optionally substituted with one, two, or three substituents selected from alkyl, alkoxy, alkylthio, halo, alkenyl, amino, monosubstituted amino, disubstituted amino, or benzofuran-2-yl, preferably a pyrimidin-2-yl ring optionally substituted with a phenyl ring optionally substituted with one, two, or three substituents selected from methyl, methoxy, methylthio, chloro, fluoro, vinyl, dimethylamino, most preferably pyrimidin-2-yl, 5-(benzofuran-2-yl)pyrimidin-2-yl, 5-phenylpyrimidin-2-yl, 5-(4-methoxyphenyl)-pyrimidin-2-yl, 5-(3,4-dimethoxyphenyl)pyrimidin-2-yl, 5-(4-dimethylamino-phenyl)pyrimidin-2-yl, 5-(4-methylphenyl)pyrimidin-2-yl, 5-(4-

methylthiophenyl)-pyrimidin-2-yl, 5-(4-vinylphenyl)pyrimidin-2-yl, or 5-(4-fluorophenyl)-pyrimidin-2-yl.

Another particularly preferred group of compounds is that wherein:

- 5 (ii) Ar^2 is a pyrazolyl ring optionally substituted with one or two substituents selected from alkyl or phenyl which is optionally substituted with a group selected from alkyl, halo or $-\text{SO}_2\text{R}$ (where R is alkyl, alkoxy, amino, mono- or disubstituted amino), preferably pyrazol-4-yl optionally substituted with one or two substituents selected from methyl or phenyl which is optionally substituted with a group selected from methyl, 2-propyl, fluoro, chloro, methoxy or $-\text{SO}_2\text{NH}_2$, more preferably 1-(4-methylphenyl)pyrazol-4-yl, 1-[4-(2-propyl)phenyl]pyrazol-4-yl, 1-(4-aminosulfonylphenyl)pyrazol-4-yl or 1-(4-fluorophenyl)pyrazol-4-yl.
- 10

(iii) A third particularly preferred group of compounds is that wherein:

- Ar^2 is:
- 15 pyridyl; preferably 4-pyridyl;
quinoxaline, preferably quinoxalin-2-yl; or
9H-1,3,4,9-tetraazafluorene, preferably 9H-1,3,4,9-tetraaza-fluor-2-ene.

- (c) A third more preferred group of compounds in group (I) is that wherein:
- 20 Z is $-\text{N}-$; and
B is $-\text{CH}_2\text{S}-$.

Within these preferred and more preferred groups, an even more preferred group of compounds is that wherein:

- 25 The stereochemistry at the C-3 carbon of the pyrrolidine ring is (S);
Q is a straight or branched alkylene chain of 1 to 3 carbon atoms, preferably $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)_2-$; and
 Ar^1 is naphthyl or a substituted phenyl ring.

Within these preferred and more preferred groups, particularly preferred group of compounds is that wherein:

Ar¹ is a phenyl ring which is optionally substituted with one, two or three substituents selected from alkyl, cyano, nitro, halo, alkylenedioxy, oxy-C2-C3-alkylene, alkoxy or phenoxy, preferably one or two substituents selected from methyl, chloro, fluoro, bromo, methylenedioxy; more preferably 3-chlorophenyl, 4-chlorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 3,4-ethylenedioxyphenyl, 2-naphthyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, 3,4-methylenedioxyphenyl, most preferably 2,3-dichlorophenyl, 3,4-dichlorophenyl or 3,4-methylenedioxyphenyl.

Within the above preferred, more preferred group and particularly preferred groups, an even more preferred group of compounds is that wherein:

(i) Ar² is pyrimidin-2-yl ring optionally substituted with alkyl or a phenyl ring which is optionally substituted with one, two, or three substituents selected from alkyl, alkoxy, alkylthio, halo, alkenyl, amino, monosubstituted amino, disubstituted amino or benzofuran-2-yl, preferably a pyrimidin-2-yl ring optionally substituted with a phenyl ring which is optionally substituted with one, two, or three substituents selected from methyl, methoxy, methylthio, chloro, fluoro, vinyl, dimethylamino, most preferably pyrimidin-2-yl, 5-(benzofuran-2-yl) pyrimidin-2-yl, 5-phenylpyrimidin-2-yl, 5-(4-methoxyphenyl)-pyrimidin-2-yl, 5-(3,4-dimethoxyphenyl)-pyrimidin-2-yl, 5-(4-dimethylaminophenyl)-pyrimidin-2-yl, 5-(4-methylphenyl)pyrimidin-2-yl, 5-(4-methylthiophenyl)pyrimidin-2-yl, 5-(4-vinylphenyl)pyrimidin-2-yl, or 5-(4-fluorophenyl)pyrimidin-2-yl.

(ii) Another particularly preferred group of compounds is that wherein:

Ar² is a pyrazolyl ring optionally substituted with one or two substituents selected from alkyl or phenyl which is optionally substituted with a group selected from alkyl, halo, or -SO₂R (where R is alkyl, alkoxy, amino, mono- or disubstituted amino), preferably pyrazol-4-yl optionally substituted with one or two substituents selected from methyl or phenyl which is optionally substituted with a group selected

from methyl, 2-propyl, fluoro, chloro, methoxy, or $-\text{SO}_2\text{NH}_2$, more preferably 1-(4-methylphenyl)-pyrazol-4-yl; 1-(4-methoxyphenyl)pyrazol-4-yl, 1-[4-(2-propyl)phenyl]pyrazol-4-yl, 1-(4-aminosulfonylphenyl)pyrazol-4-yl or 1-(4-fluorophenyl)pyrazol-4-yl.

5 (iii) A third particularly preferred group of compounds is that wherein:

Ar^2 is:

pyridyl, preferably 4-pyridyl;

quinoxaline, preferably quinoxalin-2-yl; or

9H-1,3,4,9-tetraazafluorene, preferably 9H-1,3,4,9-tetraaza-fluor-2-ene.

10

(d) A third more preferred group of compounds in group (I) is that wherein:

Z is $-\text{N}-$; and

B is $-\text{CH}_2\text{O}-$.

15

Within these preferred and more preferred groups, an even more preferred group of compounds is that wherein:

The stereochemistry at the C-3 carbon of the pyrrolidine ring is (S);

Q is a straight or branched alkylene chain of 1 to 3 carbon atoms, preferably $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)_2-$; and

20

Ar^1 is naphthyl or a substituted phenyl ring.

Within these preferred and more preferred groups, particularly preferred group of compounds is that wherein:

25

Ar^1 is a phenyl ring which is optionally substituted with one, two, or three substituents selected from alkyl, cyano, nitro, halo, alkylenedioxy, oxy-C2-C3-alkylene, alkoxy or phenoxy, preferably one or two substituents selected from methyl, chloro, fluoro, bromo, methylenedioxy; more preferably 3-chlorophenyl, 4-chlorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 3,4-ethylenedioxyphenyl, 2-naphthyl,

2,3-dichlorophenyl, 3,4-dichlorophenyl, 3,4-methylenedioxyphenyl, most preferably 2,3-dichlorophenyl, 3,4-dichlorophenyl or 3,4-methylenedioxyphenyl.

Within the above preferred, more preferred group and particularly preferred groups, an even more preferred group of compounds is that wherein:

(i) Ar^2 is pyrimidin-2-yl ring optionally substituted with alkyl or a phenyl ring which is optionally substituted with one, two, or three substituents selected from alkyl, alkoxy, alkylthio, halo, alkenyl, amino, monosubstituted amino, disubstituted amino or benzofuran-2-yl, preferably a pyrimidin-2-yl ring optionally substituted with a phenyl ring which is optionally substituted with one, two, or three substituents selected from methyl, methoxy, methylthio, chloro, fluoro, vinyl, dimethylamino, most preferably pyrimidin-2-yl, 5-(benzofuran-2-yl) pyrimidin-2-yl, 5-phenylpyrimidin-2-yl, 5-(4-methoxyphenyl)-pyrimidin-2-yl, 5-(3,4-dimethoxyphenyl)pyrimidin-2-yl, 5-(4-dimethylaminophenyl)-pyrimidin-2-yl, 5-(4-methylphenyl)pyrimidin-2-yl, 5-(4-methylthiophenyl)pyrimidin-2-yl, 5-(4-vinylphenyl)pyrimidin-2-yl, or 5-(4-fluorophenyl)pyrimidin-2-yl.

(ii) Another particularly preferred group of compounds is that wherein:
 Ar^2 is a pyrazolyl ring optionally substituted with one or two substituents selected from alkyl or phenyl which is optionally substituted with a group selected from alkyl, halo, or $-\text{SO}_2\text{R}$ (where R is alkyl, alkoxy, amino, mono- or disubstituted amino), preferably pyrazol-4-yl optionally substituted with one or two substituents selected from methyl or phenyl which is optionally substituted with a group selected from methyl, 2-propyl, fluoro, chloro, methoxy, or $-\text{SO}_2\text{NH}_2$, more preferably 1-(4-methylphenyl)-pyrazol-4-yl, 1-(4-methoxyphenyl)pyrazol-4-yl, 1-[4-(2-propyl)phenyl]pyrazol-4-yl, 1-(4-aminosulfonylphenyl)pyrazol-4-yl or 1-(4-fluorophenyl)pyrazol-4-yl.

(iii) A third particularly preferred group of compounds is that wherein:
 Ar^2 is:

pyridyl, preferably 4-pyridyl;
quinoxaline, preferably quinoxalin-2-yl; or

9H-1,3,4,9-tetraazafluorene, preferably 9H-1,3,4,9-tetraaza-fluor-2-ene.

(e) A fifth more preferred group of compounds in group (I) is that wherein:

Z is -N-; and

5 B -CH₂N-.

(II) Another preferred group of compounds is that wherein:

Z is -N-;

R¹ is hydrogen; and

10 A is -NHCONH-.

Within this preferred group (II), a more preferred group of compounds is that wherein:

B is a bond.

15

Within these preferred and more preferred groups, a particularly preferred group of compounds is that wherein:

The stereochemistry at the C-3 carbon of the pyrrolidine ring is (S);

Q is a straight or branched alkylene chain of 1 to 3 carbon atoms, preferably

20 -CH₂-, -CH(CH₃)-, -C(CH₃)₂-;

Ar¹ is a naphthyl or a phenyl ring which is optionally substituted with one, two or three substituents, selected from alkyl, cyano, nitro, halo, methylenedioxy, ethylenedioxy, alkoxy or phenoxy; preferably one or two substituents methyl, chloro, fluoro, bromo, methylenedioxy; more preferably 3-chlorophenyl, 4-chlorophenyl, 4-fluoro-3-methoxy-phenyl, 2,4-difluorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 3,4-ethylenedioxyphenyl, 2-naphthyl, 2,3-dichlorophenyl, 3,4-dichloro-phenyl, 3,4-methylenedioxyphenyl, most preferably 2,3-dichlorophenyl, 3,4-dichloro-phenyl or 3,4-methylenedioxyphenyl; and

Ar² is an aryl ring, preferably a phenyl ring which is optionally substituted with one, two, or three substituents selected from alkyl or alkoxy, preferably a phenyl ring

30

optionally substituted with one or two substituents selected from methyl or methoxy; more preferably phenyl, 3-methylphenyl, 4-methylphenyl, 3-methoxyphenyl or 4-methoxyphenyl.

5 Furthermore preferred embodiments of the present invention are:

(Z) Compounds of Formula (I) as defined as the first aspect of the present invention wherein Z is -N-; R¹ is hydrogen and A is -NHC(O)-.

10 (1) The compound of (Z) wherein Ar¹ is a naphthyl or a phenyl ring substituted with one, two, or three substituents selected from alkyl, cyano, nitro, halo, methylenedioxy, ethylenedioxy, alkoxy or phenoxy.

(2) The compound of (1) wherein Q and B are -CH₂-.

15

(3) The compound of (2) wherein Ar² is an aryl ring.

(4) The compound of (3) wherein:

20

Ar¹ is a phenyl ring substituted with one, two, or three substituents selected from methyl, chloro, fluoro, bromo, or methylenedioxy; and

Ar² is a phenyl ring optionally substituted with one, two, or three substituents selected from alkoxy, alkylthio, halo, amino, -NHC(O)R' (where R' is alkyl or optionally substituted phenyl), hydroxy, or -SO₂Me.

25

(5) The compound of (4) wherein Ar¹ is 3-chlorophenyl, 4-chlorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-ethylenedioxyphenyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, or 3,4-methylenedioxyphenyl.

- 5 (6) The compound of (5) wherein Ar² is a phenyl ring optionally substituted with one, two, or three substituents selected from methyl, methylthio, hydroxy, methoxy, acetyl, chloro, fluoro, bromo, or -NHC(O)R' (where R' is methyl or a phenyl ring optionally substituted with one, two, three, or four substituents selected from methyl, methoxy, fluoro, or chloro).
- 10 (7) The compound of (6) wherein Ar² is phenyl, 3-(2,5-difluoro-4-chlorophenyl-carbonylamino)phenyl, 4-(2,3,4,5-tetrafluorophenylcarbonylamino)phenyl, 3-(2,4-dimethoxyphenylcarbonylamino)phenyl, 3-(phenylcarbonylamino)phenyl, 3-acetylaminophenyl, 3-fluoro-4-hydroxyphenyl, 3-fluorophenyl, 3-methoxy-phenyl, or 4-hydroxy-3-methoxyphenyl.
- 15 (8) The compound of (7) wherein:
Ar¹ is 2,3-dichlorophenyl; and
Ar² is 3-fluoro-4-hydroxyphenyl;
namely, *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*S*)-ylmethyl]-2-(3-fluoro-4-hydroxyphenyl)acetamide.
- 20 (9) The compound of (7) wherein:
Ar¹ is 2,3-dichlorophenyl; and
Ar² is 3-acetylaminophenyl;
namely, *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*S*)-ylmethyl]-2-(3-acetylaminophenyl)acetamide.
- 25 (10) The compound of (1) wherein:
Q is -CH₂- and B is -(CH₂)₂;
- 30 (11) The compound of (10) wherein Ar² is a heteroaryl ring.

(12) The compound of (11) wherein:

Ar¹ is a phenyl ring substituted with one, two, or three substituents methyl, chloro, fluoro, bromo, or methylenedioxy.

5 (13) The compound of (12) wherein Ar¹ is 3-chlorophenyl, 4-chlorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-ethylenedioxyphenyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, or 3,4-methylenedioxyphenyl.

10 (14) The compound of (13) wherein Ar² is a pyrimidin-2-yl ring optionally substituted at the 5-position with a phenyl ring which is optionally substituted with one, two, or three substituents selected from methyl, methoxy, methylthio, chloro, fluoro, vinyl, or dimethylamino.

15 (15) The compound of (14) wherein Ar² is pyrimidin-2-yl, 5-phenylpyrimidin-2-yl, 5-(4-methoxyphenyl)pyrimidin-2-yl, 5-(3,4-dimethoxyphenyl)pyrimidin-2-yl, 5-(4-methylthiophenyl)pyrimidin-2-yl, 5-(4-dimethylaminophenyl)pyrimidin-2-yl, 5-(4-methylphenyl)pyrimidin-2-yl or 5-(4-fluorophenyl)pyrimidin-2-yl.

20 (16) The compound of (15) wherein:

Ar¹ is 2,3-dichlorophenyl; and

Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl;

namely, *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-ylmethyl]-3-[5-(4-methoxyphenyl)-pyrimidin-2-yl]propionamide.

25

(17) The compound of (15) wherein:

Ar¹ is 3,4-dichlorophenyl; and

Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl;

namely, *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-ylmethyl]-3-[5-(4-methoxyphenyl)-pyrimidin-2-yl]propionamide.

30

- (18) The compound of (1) wherein Q is -CH₂- and B is -CH₂S-.
- (19) The compound of (18) wherein:
Ar² is a heteroaryl ring.
- (20) The compound of (19) wherein:
Ar¹ is a phenyl ring optionally substituted with one, two, or three substituents methyl, chloro, fluoro, bromo, or methylenedioxy.
- (21) The compound of (20) wherein Ar¹ is 3-chlorophenyl, 4-chlorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-ethylenedioxyphenyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, or 3,4-methylenedioxyphenyl.
- (22) The compound of (21) wherein Ar² is a pyrimidin-2-yl ring optionally substituted at the 5-position with a phenyl ring which optionally substituted with one, two, or three substituents selected from methyl, methoxy, methylthio, chloro, fluoro, vinyl or dimethylamino.
- (23) The compound of (22) wherein Ar² is pyrimidin-2-yl, 5-phenylpyrimidin-2-yl, 5-(4-methoxyphenyl)pyrimidin-2-yl, 5-(3,4-dimethoxyphenyl)pyrimidin-2-yl, 5-(4-methylthiophenyl)pyrimidin-2-yl, 5-(4-dimethylaminophenyl)pyrimidin-2-yl, 5-(4-methylphenyl)pyrimidin-2-yl or 5-(4-fluorophenyl)pyrimidin-2-yl.
- (24) The compound of (23) wherein:
Ar¹ is 2,3-dichlorophenyl; and
Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl;
namely, *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-ylsulfanyl]acetamide.

- (25) The compound of (24) wherein:
Ar¹ is 3,4-dichlorophenyl; and
Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl;
namely, *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-[5-(4-methoxy-
phenyl)pyrimidin-2-ylsulfanyl]acetamide.
- (26) The compound of (24) wherein:
Ar¹ is 3,4-methylenedioxyphenyl; and
Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl;
namely, *N*-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-ylmethyl]-2-[5-(4-
methoxyphenyl)pyrimidin-2-ylsulfanyl]acetamide.
- (27) The compound of (1) wherein Q is -CH₂- and B is -CH₂O-.
- (28) The compound of (27) wherein:
Ar² is a heteroaryl ring.
- (29) The compound of (28) wherein:
Ar¹ is a phenyl ring optionally substituted with one, two, or three
substituents methyl, chloro, fluoro, bromo, or methylenedioxy.
- (30) The compound of (29) wherein Ar¹ is 3-chlorophenyl, 4-chlorophenyl, 3,4-
difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 4-fluoro-3-
methoxyphenyl, 2,4-difluorophenyl, 3,4-ethylenedioxyphenyl, 2,3-
dichlorophenyl, 3,4-dichlorophenyl, or 3,4-methylenedioxyphenyl.
- (31) The compound of (30) wherein Ar² is a pyrimidin-2-yl ring optionally
substituted at the 5-position with a phenyl ring which optionally substituted
with one, two, or three substituents selected from methyl, methoxy, methylthio,
chloro, fluoro, vinyl or dimethylamino.

(32) The compound of (31) wherein Ar² is pyrimidin-2-yl, 5-phenylpyrimidin-2-yl, 5-(4-methoxyphenyl)pyrimidin-2-yl, 5-(3,4-dimethoxyphenyl)pyrimidin-2-yl, 5-(4-methylthiophenyl)pyrimidin-2-yl, 5-(4-dimethylaminophenyl)pyrimidin-2-yl, 5-(4-methylphenyl)pyrimidin-2-yl or 5-(4-fluorophenyl)pyrimidin-2-yl.

5

(33) The compound of (32) wherein:

Ar¹ is 2,3-dichlorophenyl; and

Ar² is 5-(4-methoxyphenyl)pyrimidin-2-yl;

namely, *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-yloxy]acetamide.

10

(34) The compound of (32) wherein:

Ar¹ is 3,4-dichlorophenyl; and

Ar² is 5-(3,4-dimethoxyphenyl)pyrimidin-2-yl;

namely, *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-[5-(3,4-dimethoxyphenyl)pyrimidin-2-yloxy]acetamide.

15

(35) The compound of (32) wherein:

Ar¹ is 2,3-dichlorophenyl; and

Ar² is 5-(4-methylthiophenyl)pyrimidin-2-yl;

namely, *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-[5-(4-methylthiophenyl)pyrimidin-2-yloxy]acetamide.

20

(36) The compound of (Z) wherein R¹ is hydrogen and A is -NHC(O)NH-.

25

(37) The compound of (36) wherein Ar¹ is a naphthyl or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, cyano, nitro, halo, methylenedioxy, ethylenedioxy, alkoxy or phenoxy.

30

(38) The compound of (37) wherein Q is -CH₂- and B is a bond or -CH₂-.

(39) The compound of (38) wherein Ar² is an aryl ring.

(40) The compound of (39) wherein:

Ar¹ is a phenyl ring optionally substituted with one, two, or three substituents selected from methyl, chloro, fluoro, bromo, or methylenedioxy; and

Ar² is a phenyl ring optionally substituted with one or two substituents selected from alkyl or alkoxy.

(41) The compound of (40) wherein Ar¹ is 3-chlorophenyl, 4-chlorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-ethylenedioxyphenyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, or 3,4-methylenedioxyphenyl.

(42) The compound of (41) wherein Ar² is a phenyl ring optionally substituted with one or two substituents selected from methyl or methoxy.

(43) The compound of (42) wherein Ar² is phenyl, 3-methylphenyl, 4-methylphenyl, 3-methoxyphenyl or 4-methoxyphenyl.

Specifically preferred compounds of the present invention are:

N-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)-pyrimidin-2-ylsulfanyl]acetamide.

N-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-methoxyphenyl)-pyrimidin-2-yl]propionamide.

N-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-methoxyphenyl)-pyrimidin-2-yl]propionamide hydrochloride salt.

N-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)-pyrimidin-2-ylsulfanyl]acetamide.

N-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-ylsulfanyl]acetamide hydrochloride salt.

N-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(3,4-dimethoxyphenyl)pyrimidin-2-yl]propionamide.

N-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-ylsulfanyl]acetamide hydrochloride salt.

5 *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-yloxy]acetamide.

N-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-biphenyl)pyrimidin-2-yl]propionamide.

10 *N*-[1-(3,4-dichlorobenzyl)-1-methyl-3-(*RS*)-{[2-(5-phenylpyrimidin-2-ylsulfanyl)-acetyl amino]methyl}pyrrolidinium iodide.

N-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(3,4-dimethoxyphenyl)pyrimidin-2-yl]propionamide.

N-[1-(3-chlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-methoxybenzyl)pyrimidin-2-yl]propionamide.

15 *N*-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-phenylpyrimidin-2-ylsulfanyl]acetamide.

N-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(3,4-dimethoxyphenyl)pyrimidin-2-yloxy]acetamide.

20 *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methylthiophenyl)pyrimidin-2-yloxy]acetamide.

N-[1-(3-chlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-ylsulfanyl]acetamide.

N-[1-(2,3-dihydrobenzofuran-5-yl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-ylsulfanyl]acetamide.

25 *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-phenylpyrimidin-2-ylsulfanyl]acetamide.

N-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-methoxyphenyl)pyrimidin-2-yl]propionamide hydrochloride salt.

30 *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-yloxy]acetamide.

N-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-methoxyphenyl)pyrimidin-2-yl]propioamide hydrochloride salt.

N-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*S*)-ylmethyl]-2-(3-fluoro-4-hydroxyphenyl)acetamide.

5 *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-ylmethyl]-2-(3-acetylaminophenyl)acetamide.

10 The compounds of the present invention can be prepared in a number of ways known to one skilled in the art. Preferred methods include, but are not limited to, the general synthetic procedures described below.

15 The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Co., (Milwaukee, Wisconsin, USA), Bachem (Torrance, California, USA), Emka-Chemie, or Sigma (St. Louis, Missouri, USA), Maybridge (Dist: Ryan Scientific, P.O. Box 6496, Columbia, SC 92960), Bionet Research Ltd., (Cornwall PL32 9QZ, UK), Menai Organics Ltd., (Gwynedd, N. Wales, UK), Butt Park Ltd., (Dist. Interchim, Montlucon Cedex, France) or are prepared by methods known to those skilled in the art following procedures set forth in references such as *Fieser and Fieser's Reagents for Organic Synthesis*, Volumes 20 1-17 (John Wiley and Sons, 1991); *Rodd's Chemistry of Carbon Compounds*, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989), *Organic Reactions*, Volumes 1-40 (John Wiley and Sons, 1991), *March's Advanced Organic Chemistry*, (John Wiley and Sons, 1992), and *Larock's Comprehensive Organic Transformations* (VCH Publishers Inc., 1989). These schemes are merely illustrative of some methods by which 25 the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to one skilled in the art having referred to this disclosure.

30 The starting materials and the intermediates of the reaction may be isolated and purified if desired using conventional techniques, including but not limited to filtration,

distillation, crystallization, chromatography, and the like. Such materials may be characterized using conventional means, including physical constants and spectral data.

Synthesis of Compounds of Formula (I)

- 5 In general, compounds of Formula (I) where Z is -N- and A, B, R¹, R², R³, R⁴, R⁵, R⁶, Q, Ar¹ and Ar² are as defined in the first aspect of the present invention are prepared from 3-aminomethyl-pyrrolidines (Ia) or (Ib) or 3-hydroxymethylpyrrolidine (Ic) as shown in Fig. 1 below. A compound of Formula (I) where Z is -N- can be converted to a corresponding compound of Formula (I) where -N⁺R- X⁻, if desired, by alkylation with an
- 10 alkylating agent RX where R is as defined in the first aspect of the present and X is a leaving group.




Fig. 1


Synthesis of compounds of Formulae (Ia), (Ib) and (Ic) and their conversion to compounds of Formula (I) are described in detail in Schemes A-D and E-K, respectively.

Synthesis of compounds of Formulae (Ia), (Ib) and (Ic)

Synthesis of Compounds of Formula (Ia)

Scheme A


A compound of Formula (Ia) where R¹ is hydrogen is prepared as shown in Scheme A below.



Treatment of commercially available dimethyl itaconate of formula 1 with ammonia gives 4-aminocarbonyl-2-pyrrolidinone 2. Reduction with a suitable reducing agent (such as lithium aluminum hydride, diborane, and the like) in an aprotic organic solvent such as tetrahydrofuran gives 3-aminomethylpyrrolidine 3. Protection of the primary amino group as the benzylimine, followed by treatment with di-*tert*-butyl dicarbonate in the presence of a base (e.g., sodium hydroxide, sodium carbonate, triethylamine, and the like) gives 3-[(benzylideneaminomethyl)-1-*tert*-butoxycarbonylpyrrolidine which upon a mild acidic workup provides the desired 3-aminomethyl-1-*tert*-butoxycarbonylpyrrolidine of Formula (Ia). Compound (Ia) is then converted to a compound of Formula (I) as described in Schemes E-K below.

Scheme B

A compound of Formula (Ib) where R¹ is hydrogen can be prepared as described in Scheme B below.



5

Treatment of 1-benzyl-5-oxo-pyrrolidine-3-carboxylic acid methyl ester 4 with ammonia provides 3-aminocarbonyl-1-benzyl-2-pyrrolidinone 5. Reduction of 5 with a suitable reducing agent (such as lithium aluminum hydride, diborane and the like) in an aprotic organic solvent such as tetrahydrofuran gives 3-aminomethyl-1-benzylpyrrolidine 6. Protection of the primary amino group as the *tert*-butoxycarbonyl, followed by debenzylation under standard hydrogenation reaction conditions provides 3-(*tert*-butoxycarbonylamino)methylpyrrolidine 8. Compound 8 is then converted to a compound of Formula (Ib) by reacting it with a compound of formula $\text{Ar}^1\text{-Q-Y}$ where Y is an aldehyde ($-\text{CHO}$) or a keto ($-\text{C(O)R}$ where R is alkyl) group, followed by removal of the Boc group. The reaction of 8 with $\text{Ar}^1\text{-Q-Y}$ is carried out under reductive amination reaction conditions i.e., in the presence of a suitable reducing agent (e.g., sodium cyanoborohydride, sodium triacetoxyborohydride, and the like) and an organic acid (e.g.,

15


glacial acetic acid, trifluoroacetic acid, and the like) at ambient temperature. Suitable solvents for the reaction are halogenated hydrocarbons (e.g., 1,2-dichloroethane, chloroform, and the like).

5 In general, compounds of formula $\text{Ar}^1\text{-Q-Y}$ are commercially available. For example, benzaldehyde, acetophenone, 3,5-dichlorobenzaldehyde, 2-phenylpropionaldehyde, and the like are commercially available. Compound (Ib) is then converted to a compound of Formula (I) as described in Schemes E-K below.

10

Scheme C

Alternatively, a compound of Formula (Ib) where R^1 is hydrogen is prepared as shown in Scheme C below.




15

Treatment of commercially available dimethyl itaconate of formula 1 with an amine of formula 9 (where Ar^1 is as defined in the Summary of the Invention) provides a compound of formula 10 which is converted to a compound of Formula (Ib) as described above. Compounds of formula 9 such as benzylamine, 3,4-dichlorobenzylamine, phenethylamine, and the like are commercially available.

20

Preparation of Compounds of Formula (Ic)Scheme D.

A compound of Formula (Ic) where R¹ is hydrogen is prepared as shown in Scheme D below.



Reduction of 1-benzyl-5-oxo-pyrrolidine-3-carboxylic acid methyl ester **11** with a suitable reducing agent (such as lithium aluminum hydride, borane, and the like) gives N-benzyl-3-hydroxymethylpyrrolidine **12**. Removal of the benzyl group under catalytic hydrogenation reaction conditions followed by reaction of 3-hydroxymethylpyrrolidine **13** with di-*tert*-butyl dicarbonate in the presence of a base (e.g., sodium hydroxide, sodium carbonate, and the like) gives *N*-*tert*-butoxycarbonyl-3-hydroxymethylpyrrolidine of Formula (Ic). Compound (Ic) is then converted to a compound of Formula (I) as shown below.




Synthesis of Compounds of Formula (I) from Compounds of Formulae I(a-c)

Compounds of Formula (I) where Z is -N- and R¹ is hydrogen are prepared from compounds of Formulae I(a-c) as described in Schemes E - K below.

5

Scheme E

Compounds of Formula (I) where A is -N(R²)C(O)- where R² is hydrogen are prepared from a compound of Formula (Ia) or (Ib) as described in Scheme E below.

Alternate Route

10

Reaction of a compound of Formula (Ia) with a compound of formula **14** where B is as defined in the Summary of the Invention and X is a leaving group under acylating conditions such as a halo (particularly Cl or Br) or a hydroxy group gives a compound of formula **16**. The reaction conditions employed for the preparation of **16** depend on the nature of the X group. If X is a hydroxy group, the reaction is carried out in the

15

presence of a suitable coupling agent (e.g., N,N-dicyclohexylcarbodiimide, 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide, and the like). If X is a halide the reaction is carried out in the presence of a non-nucleophilic organic base (e.g., triethylamine or pyridine, preferably pyridine). Suitable organic solvent are methylene chloride, tetrahydrofuran, and the like.

Alternatively, 16 can be prepared by heating (Ia) with an acid anhydride. Suitable solvents for the reaction are tetrahydrofuran, dioxane, and the like. Compounds of formula 14 such as 4-(2,5-dimethylphenyl)-4-oxobutyric acid, 4-(acetylphenoxy)acetic acid, N-phenylsulfonylglycine, 2-(6-methoxynaphth-2-yl)-2-methylacetic acid, 3-benzenesulfonyl-propionic acid, 4-(thiophen-2-ylpyrazol-1-yl)acetic acid, 2-(1-acetylnaphth-2-yloxy)-2-methylacetic acid, 2-(4-methyl[1,2,3]thiadiazol-5-ylsulfonyl)acetic acid, 2-(quinoxalin-2-ylsulfonyl)acetic acid are commercially available.

Treatment of 16 with an aqueous acid or anhydrous acid such as hydrochloric acid or trifluoroacetic acid in dichloromethane gives a compound of formula 17 which is then converted to a compound of Formula (I) by procedures well known in the art. Some such procedures are described below.

A compound of Formula (I) can be prepared:

(i) by reacting a compound of Formula 17 with a compound of formula 18 where Y is an aldehyde (-CHO) or a keto (-C(O)R where R is alkyl) group as described previously; or

(ii) by reacting a compound of formula a compound of Formula 17 with a compound of formula 18 where Y is a leaving group under alkylating conditions such as halo (e.g., chloro, bromo or iodo) or sulfonyloxy group (e.g., methylsulfonyloxy or 4-methylphenylsulfonyloxy or trifluoromethylsulfonyloxy). The reaction is carried out in the presence of a base such as sodium carbonate, sodium hydride, triethylamine and the


like. Suitable solvents are aprotic organic solvents such as tetrahydrofuran, N,N-dimethylformamide, and the like.

In general, compounds of formula 18 where Y is an aldehyde or ketone group
5 are commercially available. For example, benzaldehyde, acetophenone, 3,5-
dichlorobenzaldehyde, 2-phenylpropionaldehyde, and the like are commercially
available. Aryl halides such as benzyl bromide, 3,4-dichlorobenzyl bromide, and
the like are also commercially available. Others can be prepared from suitable starting
materials such as phenylacetic acid, phenylpropanol, benzyloxyethanol, 3,5-
10 dichlorobenzaldehyde, 2-phenylpropionaldehyde, etc., by reducing the aldehyde, ketone
or carboxy group to an alcohol, followed by treatment with a suitable halogenating agent
(e.g., thionyl chloride, thionyl bromide, carbon tetrabromide in the presence of
triphenylphosphine, and the like) or sulfonylating agent (e.g., methylsulfonyl chloride,
para-toluenesulfonyl chloride and triflic anhydride) respectively. Suitable aldehyde,
15 ketone or carboxy reducing agents include lithium aluminum hydride, borane and the like.

Alternatively, a compound of Formula (I) can be prepared directly by reacting a
compound of Formula (Ib) with a compound of formula 14 or 15 utilizing the reaction
conditions described above.

Scheme F

Compounds of Formula (I) where Z is -N-, R¹ is hydrogen and A is -N(R²)C(O)N(R³)- or -N(R²)C(S)N(R³)- are prepared as described in Scheme F below:



5

A compound of Formula (I) where A is a urea/thiourea group can be prepared from a compound of Formula (Ia) by first preparing a compound of formula 19 either:

- (i) by reacting a compound of Formula (Ia) with an activating agent such as carbonyl diimidazole/ thiocarbonyl diimidazole, followed by nucleophilic displacement of the imidazole group with a primary or secondary amine. The reaction occurs at ambient temperature. Suitable solvents include polar organic solvents (e.g., tetrahydrofuran, dioxane and the like);
- (ii) by reacting a compound of Formula (Ia) with a carbamoyl/thiocarbamoyl halide. The reaction is carried out in the presence of a non-nucleophilic organic base. Suitable solvents for the reaction are dichloromethane, 1,2-dichloroethane, tetrahydrofuran or pyridine; or
- (iii) by reacting a compound of Formula (Ia) with an isocyanate/isothiocyanate in an aprotic organic solvent (e.g., benzene, tetrahydrofuran, dimethylformamide and the like).


15

Compound 19 is then converted to a compound of Formula (I) as described in Scheme E above.

Compound (I) can be prepared directly from a compound of Formula (IIb) by carrying out the steps (i)-(iii) above.

Scheme G

Compounds of Formula (I) where Z is -N-, R¹ is hydrogen and A is -N(R²)SO₂- where R² is hydrogen are prepared as described in Scheme G below:



10

15


A compound of Formula (I) where A is a sulfonamido group can be prepared by reacting a compound of Formula (Ia) with a sulfonyl halide, utilizing the reaction conditions described in method (ii) of Scheme D above to give a compound of formula 20 which is then converted to a compound of Formula (I) as described in Scheme E above.

Sulfonyl halides are commercially available or may be prepared by methods such as those described in (1) Langer, R. F.; *Can. J. Chem.* 61, 1583-1592, (1983); (2) Aveta, R.; et. al.; *Gazzetta Chimica Italiana*, 116, 649-652, (1986); (3) King, J. F. and Hillhouse, J. H.; *Can. J. Chem.*; 54, 498, (1976); and (4) Szymonifka, M. J. and Heck, J. V.; *Tet. Lett.*; 30, 2869-2872, (1989).

20

Scheme H

Compounds of Formula (I) where Z is -N-, R¹ is hydrogen and A is -N(R²)SO₂N(R³)- where R² is hydrogen are prepared as described in Scheme H below:



5


A compound of Formula (I) where A is a sulfamide group can be prepared by reacting a compound of Formula (Ia) with a sulfamoyl halide, utilizing the reaction conditions described in method (ii) of Scheme E above to give a compound of formula 21 which is then converted to a compound of Formula (I) as described in Scheme C above.

10

Sulfamoyl halides are commercially available or may be prepared by methods such as those described in Graf, R; *German Patent*, 931225 (1952) and Catt, J.D. and Matler, W.L.; *J. Org. Chem.*, 39, 566-568, (1974).

Scheme I

Compounds of Formula (I) where Z is -N-, R¹ is hydrogen and A is -N(R²)C(O)O- where R² is hydrogen are prepared as described in Scheme I below:




5 A compound of Formula (I) where A is a carbamate group can be prepared by first converting a compound of Formula (Ia) to a compound of formula 23 by reacting it with an activating agent such as carbonyl diimidazole, followed by nucleophilic displacement of the imidazole group with an alcohol of formula 22. The reaction occurs at ambient temperature. Suitable solvents include polar organic solvents (e.g., tetrahydrofuran, dioxane and the like). A compound of formula 24 which is then converted to a compound of Formula (I) as described in Scheme E above.

15 Alcohols of formula 22 such as benzyl alcohol, 3-benzylpropanol, and the like are commercially available.

Scheme J

Compounds of Formula (I) where Z is -N-, R¹ is hydrogen and A is -OC(O)N(R³)- are prepared as described in Scheme J below:



5

A compound of Formula (I) where A is an inverse carbamate group can be prepared by first converting a compound of Formula (Ic) to a compound of formula 25 by reacting it with an activating agent such as carbonyl diimidazole, followed by nucleophilic displacement of the imidazole group with an amine of formula 24. The reaction occurs at ambient temperature. Suitable solvents include polar organic solvents (e.g., tetrahydrofuran, dioxane and the like). A compound of formula 25 which is then converted to a compound of Formula (I) as described in Scheme E above.

10


It will be recognized by one skilled in the art that above procedures can be utilized to prepare compounds of Formula (I) directly from a compound of Formula (Ib).

15

Synthesis of a Compounds of Formula (I) where Z is -N- to a corresponding
Compound of Formula (I) where Z is -N⁺R- X⁻

Scheme K

A compound of Formula (I) where Z is -N⁺R- can be prepared from a
 5 corresponding compound of Formula (I) as shown in Scheme K below.



A compound of Formula (I) is converted to a corresponding compound of
 10 Formula (I) where Z is -N⁺R- where R is an alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxyalkyl, alkoxyalkyl, or cyanoalkyl group and X⁻ is iodide, by stirring it in neat alkyl iodide such as methyl iodide, ethyl iodide, and the like.

The iodide salt can be converted to its corresponding chloride salt by utilizing a
 15 suitable ion exchange resin such as Dowex 1x8-50.

The compounds of the invention are CCR-3 receptor antagonists and inhibit
 eosinophil recruitment by CCR-3 chemokines such as RANTES, eotaxin, MCP-2,
 MCP-3 and MCP-4. Compounds of this invention and compositions containing
 20 them are useful in the treatment of eosinophil-induced diseases such as inflammatory or allergic diseases and including respiratory allergic diseases such as asthma, allergic rhinitis, hypersensitivity lung diseases, hypersensitivity pneumonitis, eosinophilic pneumonias (e.g., chronic eosinophilic pneumonia); inflammatory bowel diseases (e.g., Crohn's disease and ulcerative colitis); and psoriasis and inflammatory dermatoses
 25 such as dermatitis and eczema.

The CCR-3 antagonistic activity of the compounds of this invention was measured by *in vitro* assays such as ligand binding and chemotaxis assays as described in more detail in Examples 14, 15 and 16. *In vivo* activity was assayed in the Ovalbumin induced Asthma in Balb/c Mice Model as described in more detail in Example 17.

In general, the compounds of this invention will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. The actual amount of the compound of this invention, i.e., the active ingredient, will depend upon numerous factors such as the severity of the disease to be treated, the age and relative health of the subject, the potency of the compound used, the route and form of administration, and other factors.

Therapeutically effective amounts of compounds of Formula (I) may range from approximately 0.01-20 mg per kilogram body weight of the recipient per day; preferably about 0.1-10 mg/kg/day. Thus, for administration to a 70 kg person, the dosage range would most preferably be about 7 mg to 0.7 g per day.

In general, compounds of this invention will be administered as pharmaceutical compositions by any one of the following routes: oral, inhalation (e.g., intranasal or oral inhalation) or parenteral (e.g., intramuscular, intravenous or subcutaneous) administration. A preferred manner of administration is oral using a convenient daily dosage regimen which can be adjusted according to the degree of affliction. Compositions can take the form of tablets, pills, capsules, semisolids, powders, sustained release formulations, solutions, suspensions, liposomes, elixirs, or any other appropriate compositions. Another preferred manner for administering compounds of this invention is inhalation. This is an effective means for delivering a therapeutic agent directly to the respiratory tract for the treatment of diseases such as asthma and other similar or related respiratory tract disorders (*see* U. S. Patent 5,607,915).

The choice of formulation depends on various factors such as the mode of drug administration and the bioavailability of the drug substance. For delivery via inhalation the compound can be formulated as liquid solutions or suspensions, aerosol propellants or dry powder and loaded into a suitable dispenser for administration. There are three types of pharmaceutical inhalation devices- nebulizer inhalers, metered-dose inhalers (MDI) and dry powder inhalers (DPI). Nebulizer devices produce a stream of high velocity air that causes the therapeutic agents (which has been formulated in a liquid form) to spray as a mist which is carried into the patient's respiratory tract. MDI's typically have the formulation packaged with a compressed gas. Upon actuation, the device discharges a measured amount of therapeutic agent by compressed gas, thus affording a reliable method of administering a set amount of agent. DPI's administer therapeutic agents in the form of a free flowing powder that can be dispersed in the patient's inspiratory air-stream during breathing by the device. In order to achieve a free flowing powder, the therapeutic agent is formulated with an excipient, such as lactose. A measured amount of the therapeutic is stored in a capsule form and is dispensed to the patient with each actuation. Recently, pharmaceutical formulations have been developed especially for drugs that show poor bioavailability based upon the principle that bioavailability can be increased by increasing the surface area i.e.; decreasing particle size. For example, U.S. Pat. No. 4,107,288 describes a pharmaceutical formulation having particles in the size range from 10 to 1,000 nm in which the active material is supported on a crosslinked matrix of macromolecules. U.S. Pat. No. 5,145,684 describes the production of a pharmaceutical formulation in which the drug substance is pulverized to nanoparticles (average particle size of 400 nm) in the presence of a surface modifier and then dispersed in a liquid medium to give a pharmaceutical formulation that exhibits remarkably high bioavailability.

The compositions are comprised of in general, a compound of Formula (I) in combination with at least one pharmaceutically acceptable excipient. Acceptable excipients are non-toxic, aid administration, and do not adversely affect the therapeutic benefit of the compound of Formula (I). Such excipient may be any solid, liquid, semi-

solid or, in the case of an aerosol composition, gaseous excipient that is generally available to one of skill in the art.

5 Solid pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk and the like. Liquid and semisolid excipients may be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, etc. Preferred liquid carriers, 10 particularly for injectable solutions, include water, saline, aqueous dextrose, and glycols.

Compressed gases may be used to disperse a compound of this invention in aerosol form. Inert gases suitable for this purpose are nitrogen, carbon dioxide, etc.

15

For liposomal formulations of the drug for parenteral or oral delivery the drug and the lipids are dissolved in a suitable organic solvent e.g. *tert*-butanol, cyclohexane (1% ethanol). The solution is lyophilized and the lipid mixture is suspended in an aqueous buffer and allowed to form a liposome. If necessary, the liposome size can be 20 reduced by sonification. (see., Frank Szoka, Jr. and Demetrios Papahadjopoulos, "Comparative Properties and Methods of Preparation of Lipid Vesicles (Liposomes)", *Ann. Rev. Biophys. Bioeng.*, 9:467-508 (1980), and D.D. Lasic, "Novel Applications of Liposomes", *Trends in Biotech.*, 16:467-608, (1998))

25

Other suitable pharmaceutical excipients and their formulations are described in *Remington's Pharmaceutical Sciences*, edited by E. W. Martin (Mack Publishing Company, 18th ed., 1990).

30


The level of the compound in a formulation can vary within the full range employed by those skilled in the art. Typically, the formulation will contain, on a

weight percent (wt%) basis, from about 0.01-99.99 wt% of a compound of Formula (I) based on the total formulation, with the balance being one or more suitable pharmaceutical excipients. Preferably, the compound is present at a level of about 1-80 wt%. Representative pharmaceutical formulations containing a compound of Formula (I) are described in Example 13.

EXAMPLES

Example 1

Synthesis of 3-(*RS*)-aminomethyl-1-*tert*-butoxycarbonylpyrrolidine



Step 1

Melted dimethyl itaconate (150 g, 0.95 mol) was added to an anhydrous solution of ammonia in methanol (7 M, 1000 mL, 7 mol) at room temperature. The reaction vessel was sealed with parafilm and vented with a needle. The solution was left to stand for 3 days, after which the solid was filtered and washed with cold methanol and dried under vacuum to yield pure 4-aminocarbonylpyrrolidin-2-one (150 g).

Step 2


To a stirred slurry of 4-aminocarbonylpyrrolidin-2-one (20 g, 156 mmol) in (100 mL) of cold (0 °C) dry tetrahydrofuran, in a 1L flask under N₂, was added a solution of lithium aluminum hydride (12 g, 0.32 mmol) in dry tetrahydrofuran (350 mL) dropwise. Once addition was complete and hydrogen evolution had abated, the cooling bath was removed and the reaction mixture was heated at reflux temperature. After the reaction was complete, the reaction mixture was allowed to cool to room

temperature and ether (300 mL) was added. Sufficient saturated Na_2SO_4 was added dropwise over 1 h to destroy excess hydride. Excess water was avoided so as to prevent two phases forming, as the diamine is highly water-soluble. The suspension was filtered over celite with copious washings with tetrahydrofuran and 15% methanol in tetrahydrofuran. The combined washings were concentrated on the rotovap and the pungent residue distilled under vacuum to yield pure 3-(*RS*)-aminomethylpyrrolidine as a colorless mobile oil (7.6 g).

Step 3

10 Benzaldehyde (5.3 g, 50 mmol) was added to a solution of 3-aminomethylpyrrolidine (5.0 g, 50 mmol) in toluene (anhydrous, 100 mL) in a 250 mL flask at room temperature under N_2 . A Dean-Stark apparatus and condenser were fitted and the reaction mixture was heated at reflux for 4 h. The reaction mixture was cooled to room temperature and di-*tert*-butyldicarbonate (5.3 g, 50 mmol) was added portion-
15 wise and the resulting solution is stirred at room temperature overnight. The reaction mixture was concentrated on the rotavap and the residue was diluted with 1 M NaHSO_4 (80 mL, 80 mmol, 1.6 equiv.) and stirred vigorously for 2 h. The reaction mixture was washed with ether to remove unwanted organic byproducts, then basified with 1 M NaOH to pH 7. Further extraction with either ether or ethyl acetate removed
20 additional unwanted byproducts. The aqueous solution was then made strongly basic (pH 12) with 1 M NaOH and extracted with ethyl acetate. The organic layer was washed with brine, dried with Na_2SO_4 , then concentrated to give the 3-(*RS*)-aminomethyl-1-*tert*-butoxycarbonyl-pyrrolidine as a colorless oil (80-90 %), which was used directly without further purification.

25

Example 2**Synthesis of 3-(*S*)-aminomethyl-1-*tert*-butoxycarbonylpyrrolidine****Step 1**

(*S*)-pyrrolidinol (30g, 0.34 mol) was dissolved in a mixture of water/dioxane (400 mL/300 mL) and *N,N*-diisopropylethylamine (120 mL, 0.69 mol, 2 equiv.) was added. This reaction mixture was cooled to 0°C and di-*tert*-butyldicarbonate (90 g, 0.41 mol, 1.2 equiv.) in solution in dioxane (100 mL) was added slowly. The resulting mixture was stirred at 0°C for 1 h then at room temperature for 8 h. The reaction mixture was then partitioned between diethyl ether and water. The aqueous phase was acidified to pH 3.0 with 1 M HCl and extracted with ethyl acetate. The organic phases were combined and washed with a brine solution. The resulting organic layer was dried over magnesium sulfate, filtered and concentrated to give, after purification by flash chromatography (methanol/CH₂Cl₂: 15/85) (*S*)-1-(*tert*-butoxycarbonylamino)-3-hydroxypyrrolidine as a white solid (40 g).

Step 2

(*S*)-1-(*tert*-Butoxycarbonylamino)-3-hydroxypyrrolidine (40 g, 0.21 mol) was dissolved in anhydrous methylene chloride (400 mL), in the presence of *N,N*-diisopropyl-ethylamine (73 mL, 0.42 mol, 2 equiv.) and cooled using an ice bath. This solution was treated with methanesulfonyl chloride (18 mL, 0.23 mol, 1.1 equiv.) over a 30 min. period and was then allowed to stir at room temperature for 4 h. The solution was concentrated and the residue dissolved in ethyl acetate (250 mL). The resulting organic solution washed with 5 % NaHCO₃ and then with brine. The organic extract was dried over magnesium sulfate, filtered and concentrated to give (*S*)-1-(*tert*-

butoxycarbonyl-amino)-3-methanesulfonyloxypyrrolidine as a dark oil (39 g) which was used directly without further purification.

Step 3


5 (S)-1-(*tert*-Butoxycarbonylamino)-3-methanesulfonyloxypyrrolidine (39 g, 0.15 mol) was dissolved in acetonitrile (250 mL) and tetra-*n*-butylammonium cyanide (75 g, 0.28 mol, 1.9 equiv.) was added. The resulting mixture was heated at 65°C for 6 h and then cooled to room temperature. Saturated NaHCO₃ (500 mL) was added and the reaction mixture extracted with toluene (750 mL and 300 mL). The combined organic
10 layers were washed with water and concentrated under reduced pressure to give a brown oil. This crude material was purified by flash chromatography with 20 % ethyl acetate in hexane to give (S)-1-(*tert*-butoxycarbonylamino)-3-cyanopyrrolidine as a yellow oil (13.5 g).

15 Step 4.

A solution of (S)-1-(*tert*-butoxycarbonylamino)-3-cyanopyrrolidine (13.5g, 67 mmol) in 50 mL of 3 % NH₄OH/methanol and Raney Nickel (0.5 g, 50 % slurry in water) was pressurized to 40 psi under H₂ atmosphere. After stirring for 12 h at room temperature, the reaction mixture was filtered through a pad of Celite and the residue
20 was washed with 100 mL of methanol. The filtrate was concentrated to dryness to give an oil (S)-1-(*tert*-butoxycarbonylamino)-3-aminomethylpyrrolidine (10 g).

Proceeding as described above, but substituting (S)-pyrrolidinol (1) with (R)-pyrrolidinol (1) gave (R)-1-(*tert*-butoxycarbonylamino)-3-aminomethylpyrrolidine.

25

Example 3**Synthesis of 3-(*RS*)-aminomethyl-1-(3,4-dichlorobenzyl)pyrrolidine****5 Step 1**


Neat 3,4-dichlorobenzylamine (18 g, 114 mmol) was added to a solution of dimethyl itaconate (20 g, 114 mmol) in methanol (200 mL) at room temperature. The solution was stirred at for 48 h then concentrated *in vacuo*. The resulting solid was divided equally into two, one portion being treated with methanolic ammonia (7 M, 300 mL, 2.1 mol). The solution was vented with a needle and allowed to stand for 2 days. The slurry of solvent and product was concentrated further and filtered. The filter cake was washed with cold methanol to give pure 3-aminocarbonyl-1-(3,4-dichlorobenzyl)-pyrrolidin-2-one (18.5 g).

15 Step 2

A suspension of 3-aminocarbonyl-1-(3,4-dichlorobenzyl)pyrrolidin-2-one in dry tetrahydrofuran (150 mL) was added slowly to a solution of lithium aluminum hydride (4.9 g, 128 mmol, 2 equiv.) in tetrahydrofuran (100 mL) under N₂ at room temperature. The reaction mixture was heated at reflux overnight, diluted with ether, and quenched with brine. After 1 h of rapid stirring, the gray mixture was filtered through celite (ethyl acetate washings) and the filtrate was concentrated. Flash chromatography with CHCl₃/MeOH/NH₃ solution (200:25:1;) gave 3-(*RS*)-aminomethyl-1-(3,4-dichloro-benzyl)pyrrolidine compound as a colorless oil (6.2 g) which was >95 % pure.

Example 4

Synthesis of *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-methoxyphenyl)pyrimidin-2-yl]propionamide hydrochloride salt

**Step 1**

Methyl 1-benzyl-5-oxo-3-pyrrolidinecarboxylate (50g, 0.21 mol) was stirred in a solution of ammonia in methanol (7 M, 400 mL, 2.8 mol). After 2 days, the solution was concentrated with heating (50 °C) to about 300 mL at which point all solids remained dissolved. The reaction was then allowed to cool down and the resulting solid was filtered, washed with ether and dried to give 1-benzyl-5-oxo-pyrrolidine-3-carboxylic acid amide as colorless crystals (40 g).

Step 2

1-Benzyl-5-oxo-pyrrolidine-3-carboxylic acid amide (22 g, 0.1 mol) was added portionwise to a stirred solution of lithium aluminum hydride (9.5 g, 0.25 mol, 2.5 equiv.) in dry tetrahydrofuran (600 mL). After the initial effervescence had subsided, the reaction mixture was heated at reflux at room temperature for 24 h, at which time analysis of the reaction mixture by LCMS showed there was no starting material. The reaction mixture was quenched by dropwise addition of saturated sodium sulphate solution with stirring until no further effervescence was observed. The suspension was filtered through a celite plug, eluting with diethyl ether (200 mL). The solvent was removed to afford 3-(*RS*)-aminomethyl-1-benzylpyrrolidine as an oil (14.7 g) which was used directly without further purification.

Step 3

3-(*RS*)-Aminomethyl-1-benzylpyrrolidine (14.7 g, 80 mmol) was dissolved in methylene chloride (400 mL) with stirring. A solution of di-*tert*-butyl dicarbonate (16.8 g, 80 mmol, 1.0 equiv.) in methylene chloride (50 mL) was added dropwise and the reaction mixture was stirred overnight. The reaction mixture was concentrated and the residue was dissolved in diethyl ether (400 mL) and extracted with NaHSO₄ solution (1 M, 120 mL). The aqueous layer was basified to pH 12 with 1 M sodium hydroxide and extracted with ethyl acetate. The organic fractions were combined, washed with brine, dried over magnesium sulfate, and filtered. The organics were evaporated to afford 1-benzyl-3-(*RS*)-(N-*tert*-butoxycarbonylamino)methylpyrrolidine (22.5 g) which was used directly.

Step 4

1-Benzyl-3-(*RS*)-(N-*tert*-butoxycarbonylamino)methylpyrrolidine (22.2 g, 0.76 mol) was dissolved in a mixture of methanol and acetic acid (1:1, 100 mL) and added to a Parr flask charged with palladium on carbon (10 %, 4 g) suspended in methanol and acetic acid (1:1, 100 mL). The flask was transferred to the Parr reduction apparatus and the suspension shaken under an atmosphere of hydrogen at 60 psi for 2 days. The reaction mixture was filtered through a celite plug, eluting with a mixture of methanol/methylene chloride. The solvent was concentrated and the residue co-evaporated with toluene to afford 3-(*RS*)-(N-*tert*-butoxycarbonylamino)methylpyrrolidine as a colorless oil which was used directly.

Step 5

Sodium triacetoxyborohydride (3.2 g, 15 mmol, 1.5 equiv.) was added in one portion to a stirred solution of 3-(*RS*)-(N-*tert*-butoxycarbonylamino)methylpyrrolidine (2.0 g, 10 mmol) and 2,3-dichlorobenzaldehyde (1.9 g, 11 mmol, 1.1 equiv.) in dichloroethane (60 mL) at room temperature. The suspension was stirred overnight, then concentrated *in vacuo*. The residue was diluted with ether and quenched with 1 M hydrochloric acid. The aqueous phase was basified with 4 M sodium hydroxide to pH

12, then extracted with ethyl acetate. The organic extracts were combined and washed with brine, then dried over sodium sulfate and concentrated. Flash chromatography of the residue afforded 3-(*RS*)-(N-*tert*-butoxycarbonylaminomethyl)-1-(2,3-dichlorobenzyl)-pyrrolidine as a colorless oil (2.9 g). The oil was taken up into methylene chloride (30 mL) and treated with neat trifluoroacetic acid (5 mL). After 1 h, the volatile components were removed on a vacuum pump, then concentrated further under high vacuum to give 3-(*RS*)-aminomethyl-1-(2,3-dichlorobenzyl)pyrrolidine which was used directly without further purification.

10 Step 6

A solution of sodium ethoxide (21% wt/vol in ethanol, 10 mL, 30 mmol, 3 equiv.) was added in one portion to a suspension of 2-(4-methoxyphenyl)trimethinium perchlorate (3.3 g, 9.8 mmol) [see, Jutz, C.; Kirchlechner, R.; Seidel, H. *Chem. Ber.* 102, 2301, (1969)] and 4-amidinobutanoic acid mono-HCl (1.5 g, 9.8 mmol) [see, 15 McElvain, S.M.; Schroeder, J.P. *J. Am. Chem. Soc.*, 71, 40, (1949)] in absolute ethanol (40 mL). The reaction mixture was heated at reflux for 12 h, then cooled to room temperature. The suspension was concentrated, diluted with water, and washed with ether. The aqueous phase was then made acidic with citric acid (10 g). The precipitates were filtered, washed with water and ether, and dried under high vacuum 20 to yield 3-[4-(4-methoxyphenyl)-pyrimidin-2-yl]propanoic acid (1.85 g) as an off-white solid.


Step 7

1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.29 g, 1.5 mmol, 1.5 equiv.) was added to a suspension of 3-[4-(4-methoxyphenyl)pyrimidin-2-yl]-propanoic acid (0.26 g, 1.0 mmol, 1.0 equiv.), 3-(*RS*)-aminomethyl-1-(2,3-dichlorobenzyl)-pyrrolidine (0.26 g, 1.0 mmol), 1-hydroxybenzotriazole hydrate (0.20 g, 1.5 mmol, 1.5 equiv.), and diisopropylethylamine (0.44 mL, 2.5 mmol, 2.5 equiv.) in chloroform (2 mL). After stirring overnight at room temperature, the reaction mixture 30 was diluted with ethyl acetate, washed sequentially with 1 M sodium hydroxide, water,

1 M hydrochloric acid, water and brine, then dried over sodium sulfate and concentrated. The residue was purified by flash chromatography (0→10% methanol in methylene chloride). Clean fractions containing the desired product were combined and concentrated, diluted with dioxane and treated with 4 M HCl (0.2 mL) in dioxane. The solution was concentrated under high vacuum to give *N*-(1-(2,3-dichlorobenzyl)-pyrrolidin-3-(*RS*)-ylmethyl)-3-(5-(4-methoxyphenyl)pyrimidin-2-yl)propionamide as the HCl salt (185 mg).

Example 5

10 Synthesis of *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-nitrophenyl)pyrimidin-2-yl]propionamide



Step 1

15 Phosphorus oxychloride (83 mL, 0.79 mol) was added slowly to cold, dry, dimethyl formamide (100 mL) under N₂ at such a rate that the temperature did not rise above 5 °C. After addition was complete, 4-nitrophenylacetic acid (48 g, 0.26 mol) was added in one portion and the reaction mixture was heated to 85 °C over 1 h. After 1 h, the reaction mixture was cooled, then poured over ice. Solid sodium perchlorate monohydrate (37 g, 0.26 mol) was added to initiate precipitation of the product as the perchlorate salt. Filtration of the solid, followed by washings with cold water, 20 methanol and ether, gave 2-(4-nitrophenyl)trimethinium perchlorate (81.9 g) as a pale yellow solid.

Step 2


A solution of sodium ethoxide (21% wt/vol, in ethanol, 60 mL, 180 mmol, 3 equiv.) was added in one portion to a suspension of 2-(4-nitrophenyl)trimethinium perchlorate (20.8 g, 60 mmol) and 4-aminopropionic acid mono hydrochloride salt (9.1 g, 60 mmol) in ethanol (300 mL). The suspension was heated at room temperature overnight. The resulting suspension was filtered, washed with ethanol, cold HCl, water and ether, then dried under high vacuum to give 3-[5-(4-nitrophenyl)pyrimidin-2-yl]-propanoic acid (13.7 g) as a beige solid.

Step 3

1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (5.8 g, 30 mmol, 1.5 equiv.) was added to a suspension of 3-[4-(4-nitrophenyl)pyrimidin-2-yl]propanoic acid (6.6 g, 24 mmol, 1.2 equiv.), (1-*tert*-butoxycarbonyl)-3-(*RS*)-aminomethylpyrrolidine (4.0 g, 20 mmol), 1-hydroxybenzotriazole hydrate (4.1 g, 30 mmol, 1.5 equiv.), and triethylamine (7.0 mL, 50 mmol, 2.5 equiv.) in chloroform (50 mL). Dimethylformamide (100 mL) was added to fully dissolve the reaction components and the solution was stirred at room temperature overnight. The reaction mixture was diluted with ethyl acetate, washed sequentially with 1 M sodium hydroxide, water, 1 M hydrochloric acid, water and brine, then dried sodium sulfate and concentrated to afford a light brown solid. This material was washed with ether to afford clean *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-nitrophenyl)pyrimidin-2-yl]propionamide (4.9 g).

Example 6

Synthesis of *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-aminophenyl)pyrimidin-2-yl]propionamide



5


Hydrogen gas (balloon pressure) was introduced into a vessel containing Pd / C (10 %, 100 mg, 0.1 mmol, 0.05 equiv.) in a solution of *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-nitrophenyl)pyrimidin-2-yl]propionamide (1.0 g, 2.2 mmol) in methanol (30 mL). After 1.5 h, the reaction was terminated by purging with N₂ and the reaction mixture was filtered through celite® and concentrated. The yellow residue was dissolved in ethyl acetate, then taken into the aqueous phase using 1 M hydrochloric acid. The aqueous phase was basified to pH 11, then extracted thoroughly with ethyl acetate. The organic fractions were combined and washed with brine, then dried sodium sulfate and concentrated to afford *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[5-(4-aminophenyl)pyrimidin-2-yl]propionamide as a yellow crystalline solid (630 mg).

10

15

Example 7

Synthesis of *N*-(1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl)-2-[5-(4-methoxyphenyl)-pyrimidin-2-ylsulfanyl]acetamide

**Step 1**

Benzaldehyde (5.3 g, 50 mmol) was added to a solution of 3-(*RS*)-aminomethyl-pyrrolidine (5.0 g, 50 mmol) in toluene (anhydrous, 100 mL) in a 250 mL flask at room temperature under N₂. A Dean-Stark apparatus and condenser were fitted, the reaction vessel well lagged and the reaction mixture heated at a strong reflux for 4 h. The reaction mixture was cooled to room temperature and di-*tert*-butyl dicarbonate (5.3 g, 50 mmol) was added portion-wise and the resulting solution was stirred at room temperature overnight. The reaction mixture was concentrated on the rotavap then the residue was diluted with 1 M NaHSO₄ (80 mL, 80 mmol, 1.6 equiv.) and stirred vigorously for 2 h. The reaction mixture was washed with ether to remove unwanted organic byproducts, then basified with 1 M NaOH to pH 7. Further extraction with ethyl acetate removed additional unwanted byproducts. The aqueous solution was then made strongly basic (pH 12) with 1 M NaOH and extracted thoroughly with ethyl acetate. The organic layer was washed with brine, dried with sodium sulfate, then concentrated to give 3-(*RS*)-aminomethyl-1-*tert*-butoxycarbonylpyrrolidine as a colorless oil which was used directly without further purification.

Step 2

3-(*RS*)-aminomethyl-1-*tert*-butoxycarbonylpyrrolidine (6.0 g, 30 mmol) in methylene chloride (15 mL) was added dropwise to a cold (0 °C) solution of

chloroacetylchloride (2.4 mL, 30 mmol) and diisopropylethylamine (5.5 mL, 32 mmol) in methylene chloride under N₂. After 1 h, an additional 0.5 equiv. of chloroacetyl chloride and diisopropylamine were added. After being left at 0 °C overnight, the reaction mixture was diluted with ethyl acetate and washed briefly with water and brine, then dried over sodium sulfate and concentrated to give *N*-(1-*tert*-butoxycarbonylpyrrolidin-3-(*RS*)-ylmethyl)-2-chloroacetamide as a dark brown oil which was used without further purification.

Step 3

A solution of sodium ethoxide (21% wt/vol, 15 mL, 45 mmol, 1.5 equiv.) was added to a suspension of 2-(4-methoxyphenyl)trimethinium perchlorate (10.0 g, 30 mmol) and thiourea (3.0 g, 40 mmol, 1.3 equiv.) in ethanol (200 mL) (*see*, Krecmerova, M.; Hrebabecky, H.; Masojdkova, M.; Holy, A. *Collect. Czech. Chem. Commun.*, 61, 458, (1996)]. The reaction mixture was heated at 60 °C for 2 h. Additional quantities of thiourea and sodium ethoxide were added to the starting amounts and the reaction mixture was heated at 60 °C for an additional 1 h. The yellow suspension was cooled to room temperature, quenched with acetic acid (10 mL) and filtered. The solid was washed with water and ethanol and dried under high vacuum to give 5-(4-methoxyphenyl)-1H-pyrimidine-2-thione as a free flowing yellow powder (6.0 g).

Step 4

5-(4-Methoxyphenyl)-1H-pyrimidine-2-thione (0.70 g, 3 mmol) was added to a solution of the *N*-(1-*tert*-butoxycarbonylpyrrolidin-3-(*RS*)-ylmethyl)-2-chloroacetamide (1.15 g, ~80 % purity, ~3 mmol) and diisopropylethylamine (0.9 mL, 5 mmol, 1.5 equiv.) in dry acetonitrile. An ethanolic solution of sodium ethoxide (2.7 M, 1.2 mL, 3.2 mmol) and dimethylformamide (20 mL) were added in order to dissolve the starting mercaptan. The resulting dark brown solution was stirred at room temperature for 2 h. The reaction mixture was concentrated and the residue was chromatographed on silica gel (50-100% ethyl acetate in hexane). Fractions containing

the major product were combined and concentrated to an oily residue. Trituration of the oil with ether afforded pure *N*-(1-*tert*-butoxycarbonylpyrrolidin-3-(*RS*)-ylmethyl)-2-[5-(4-methoxyphenyl)pyrimidin-2-ylthio]-acetamide (0.60 g, 44 %) as a pale yellow solid.

5

Step 5

Neat anhydrous trifluoroacetic acid (3 mL) was added to a solution of *N*-(1-*tert*-butoxycarbonylpyrrolidin-3-(*RS*)-ylmethyl)-2-[5-(4-methoxyphenyl)pyrimidin-2-ylthio]-acetamide (0.60 g, 1.3 mmol) in methylene chloride (20 mL) at room temperature. Gas evolution was apparent immediately upon addition of the acid. After 30 min., the reaction mixture was concentrated using a Teflon dryvac system, and was then further concentrated under high vacuum to give *N*-(pyrrolidin-3-(*RS*)-ylmethyl)-2-[5-(4-methoxyphenyl)pyrimidin-2-ylthio]acetamide which was dissolved in 10 mL dichloroethane (0.13 mmol/mL).

15

Step 6

2,3-Dichlorobenzaldehyde (115 mg, 0.66 mmol, 1.5 equiv.) was added to *N*-(pyrrolidin-3-(*RS*)-ylmethyl)-2-[5-(4-methoxyphenyl)pyrimidin-2-ylthio]acetamide in dichloroethane (3.4 mL, 0.44 mmol). Excess Na(OAc)₃BH (0.2 g, 0.9 mmol, ~3 equiv.) was added and the resulting suspension was stirred vigorously overnight. The reaction mixture was diluted with ether and quenched with 1 M hydrochloric acid to afford a cloudy mixture. The organic layer was carefully removed and the aqueous phase made basic with 5 M aqueous sodium hydroxide to pH 11. After thorough extraction with ethyl acetate, the combined organic phases was washed with brine, then dried over sodium sulfate and concentrated. Chromatography on silica (0-10% methanol in ethyl acetate) afforded *N*-(1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl)-2-[5-(4-methoxyphenyl)-pyrimidin-2-ylsulfanyl]acetamide that was converted to the hydrochloride salt using 4 M hydrochloric acid solution in dioxane and ether (60 mg).

25

Proceeding as described in Step 6 above, but substituting 2,3-dichlorobenzylaldehyde (115 mg, 0.66 mmol, 1.5 equiv.) with 3,4-dichlorobenzaldehyde gave *N*-(1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl)-2-[5-(4-methoxyphenyl)pyrimidin-2-ylsulfanyl]acetamide hydrochloride salt.


5

Proceeding as described in Step 6 above, but substituting 2,3-dichlorobenzylaldehyde with 3,4-methylenedioxybenzaldehyde gave *N*-(1-(3,4-methylenedioxybenzyl)pyrrolidin-3-(*RS*)-ylmethyl)-2-[5-(4-methoxyphenyl)pyrimidin-2-ylsulfanyl]-acetamide hydrochloride salt.

10

Example 8

Synthesis of *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-yloxy]acetamide



15

Step 1

The procedure for the synthesis of 5-bromo-2-hydroxypyrimidine used is a variation of that published by Crosby and Berthold, *J. Chem. Soc.* 25, 1916, (1960). To a solution of 2-hydroxypyrimidine hydrochloride (100 g, 0.75 mol) in water (1.2 L) was added bromine (135 g, 0.84 mol) slowly with stirring. The reaction mixture was continuously stirred for about 30 min., until the red color of the solution became lighter. The solution was heated to 80 °C to drive off excess Br₂ and HBr. The solvent was concentrated further under vacuum and the residue recrystallized from 90% aqueous ethanol to afford 5-bromo-2-hydroxypyrimidine (111 g).

25

Step 2

Phosphorus oxychloride (225 mL, 2.4 mol, 1.4 equiv.) was added to a mixture of 5-bromo-2-hydroxypyrimidine (30 g, 0.17 mol) and dimethylaniline (7.5 mL) and the solution was heated at reflux under N₂ for 4 h. The dark brown reaction mixture
5 was cooled, poured over ice and extracted with ether. The organic phase was washed with bicarbonate solution, dried sodium sulfate and concentrated to afford 5-bromo-3-chloropyrimidine (25 g, 75 %) [see, Goodby, J.W.; Hird, M.; Lewis, R.A.; Toyne, K.J. *J. Chem. Soc., Chem. Commun.*, 2719, (1996)].

10 Step 3

The synthesis of (5-bromopyrimidin-2-yloxy)acetic acid followed the protocol described for similar analogues by Coppola, G.M.; Hardtmann, G.E.; Huegi, B.S. *J. Heterocycl. Chem.*, 17, 1479, (1980). Sodium hydride (5.0 g, 60 % dispersion in mineral oil, 124 mmol, 1.8 equiv.) was washed twice with dry hexane under N₂, then
15 added portionwise to a solution of methyl glycolate (9.4 g, 103 mmol, 1.5 equiv.) in toluene (150 mL). The reaction mixture was stirred at room temperature for 30 min., then 5-bromo-3-chloropyrimidine (13.3 g, 69 mmol) in toluene (50 mL) was added. The reaction mixture was heated at 60 °C overnight and concentrated. The residue was stirred rapidly with 1 M aqueous sodium hydroxide (excess) for 30 min., washed with
20 ether, then acidified to with 4 M hydrochloric acid. The resulting precipitate was collected and washed with cold water. The filtrate was extracted further with ethyl acetate, and the organic phase washed with brine, then dried sodium sulfate and concentrated to give (5-bromopyrimidin-2-yloxy)acetic acid (10.3 g).

25 Step 4

To a solution of 3-(RS)-aminomethyl-1-*tert*-butoxycarbonylpyrrolidine in dichloroethane (17 mL) were sequentially added diisopropylethylamine (5.1 mL), a solution of 2-(5-bromopyrimidin-2-yloxy)acetic acid (2.9 g, 12.2 mmol, 1.1 equiv.) in tetrahydrofuran (43 mL), 1-hydroxybenzotriazole hydrate (2.4 g, 17.5 mmol, 1.5
30 equiv.) and 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride (3.4 g,

17.6 mmol, 1.5 equiv.). The solution was stirred under N₂ overnight. The reaction mixture was diluted with ethyl acetate, washed with 1 M aqueous sodium hydroxide, 1 M hydrochloric acid, water and brine, then dried over sodium sulfate and concentrated. The residue was subjected to flash chromatography to afford clean *N*-(1-*tert*-
5 butoxycarbonylpyrrolidin-3-(*RS*)-ylmethyl)-2-(5-bromopyrimidin-2-yloxy)acetamide (4.2 g).

Step 5

4-Methoxyphenylboronic acid (1.0 g, 6.8 mmol, 1.05 equiv.) was added to a
10 solution of *N*-(1-*tert*-butoxycarbonylpyrrolidin-3-(*RS*)-ylmethyl)-2-(5-bromopyrimidin-2-yloxy)acetamide (2.7 g, 6.5 mmol) in 1-propanol (30 mL). The suspension was stirred until all ingredients had dissolved (~10 min). The resulting solution was treated with palladium acetate (29 mg, 0.13 mmol, 0.02 equiv.), triphenylphosphine (103 mg, 0.39 mmol, 0.06 equiv.), 2 M aqueous sodium carbonate (3.9 mL, 7.8 mmol, 1.2
15 equiv.) and deionized water (9 mL). The reaction mixture was heated at reflux under N₂ for 1 h. Water (20 mL) was added and the N₂ inlet was removed. After stirring at room temperature overnight, the reaction mixture was thoroughly extracted with ethyl acetate. The combined organic phase was washed with saturated sodium bicarbonate solution and brine. The organic phase was stirred with activated charcoal for 15 min,
20 dried over sodium sulfate and concentrated. Recrystallization of the residue from ethyl acetate in hexane afforded clean *N*-(1-*tert*-butoxycarbonylpyrrolidin-3-(*RS*)-ylmethyl)-2-(5-(4-methoxyphenyl)pyrimidin-2-yloxy)acetamide (2.1 g). The solid was dissolved into methylene chloride (30 mL) and neat trifluoroacetic acid (5 mL) added dropwise. After about 1 h, the solution was concentrated to an oil on a Teflon dryvac, then under
25 high vacuum to give *N*-(pyrrolidin-3-(*RS*)-ylmethyl)-2-[5-(4-methoxyphenyl)pyrimidin-2-yloxy]acetamide.


Step 6

A stock solution containing *N*-(pyrrolidin-3-(*RS*)-ylmethyl)-2-(5-(4-methoxy-
30 phenyl)pyrimidin-2-yloxy)acetamide (0.44 mmol) in dichloroethane (10 mL) was

added to a solution of 3,4-dichlorobenzaldehyde (85 mg, 0.48 mmol, 1.1 equiv.) and diisopropylethylamine (0.35 mL, 2.0 mmol, 4.5 equiv.) in dichloroethane (5 mL). Excess $\text{Na}(\text{OAc})_3\text{BH}$ (140 mg, 0.66 mmol, 1.5 equiv.) was added and the reaction mixture was stirred rapidly at room temperature overnight. The reaction mixture was quenched with methanol, concentrated and diluted with 1:1 methanol/dimethyl sulfoxide. The solution was then purified directly using preparative reverse phase chromatography to give, after treatment with 4 M hydrochloric acid in dioxane (1 mL) and concentration, *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-yloxy]-acetamide HCl salt (96 mg).

Example 9

Synthesis of *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-ylamino]acetamide hydrochloride salt



Step 1

An ethanolic solution of sodium ethoxide (2.7 M, 3.8 mL, 10 mmol, 2.9 equiv.) was added to a suspension of 2-(4-methoxyphenyl)trimethinium perchlorate salt (1.1 g, 3.4 mmol) and guanidineacetic acid (0.48 g, 4.0 mmol, 1.2 equiv.) in dehydrated ethanol (20 mL). The reaction mixture was stirred at room temperature for 30 min., then at reflux temperature for 3 h. After cooling, the sodium salt was filtered, and the cake dissolved in 20 mL of water then acidified with 1 M hydrochloric acid. The aqueous layer was extracted with ethyl acetate and the combined organic phases was washed with brine, and dried over sodium sulfate. Removal of the solvent under vacuum afforded a solid which contained ~1:1 mixture of regioisomers. The two components were separated using reversed-phase chromatography to give 2-[5-(4-

methoxyphenyl)pyrimidin-2-ylamino]-acetic acid (150 mg) and the regioisomer (100 mg).


Step 2

5 Solid 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (14 mg, 0.08 mmol, 1.5 equiv.) was added to a solution of 3-(*RS*)-aminomethyl-1-(2,3-dichlorobenzyl)-pyrrolidine (13 mg, 0.05 mmol), 2-[5-(3-methoxyphenyl)pyrimidin-2-ylamino]acetic acid (16 mg, 0.06 mmol, 1.2 equiv.), 1-hydroxybenzotriazole hydrate (10 mg, 0.08 mmol, 1.5 equiv.) and diisopropylethylamine (22 mL, 0.13 mmol, 2.5
10 equiv.) in dimethyl formamide (0.5 mL). The reaction mixture was shaken at room temperature overnight. The reaction mixture was quenched with methanol (0.3 mL), then purified directly by reverse phase preparative HPLC to give *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[5-(4-methoxyphenyl)pyrimidin-2-ylamino]acetamide HCl salt (11 mg).

15

Example 10

Synthesis of *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-4-(2,5-dimethylphenyl)-4-oxo-butyramide trifluoroacetate salt



20

To a solution (1 mL) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (55 mg, 0.28 mmol, 1.8 equiv.), 1-hydroxybenzotriazole hydrate (38 mg, 0.28 mmol, 1.8 equiv.) and triethylamine (43 mL, 0.31 mmol, 2.0 equiv.) in chloroform were added solid 4-(2,5-dimethylphenyl)-4-oxobutyric acid (39 mg, 0.19 mmol, 1.2 equiv.) and 3-(*RS*)-aminomethyl-1-(3,4-dichlorobenzyl)pyrrolidine (40 mg, 0.16 mmol) in chloroform (1 mL). The reaction mixture was shaken overnight, then
25

concentrated and diluted with dimethylsulfoxide/methanol (1:1, 1 mL). The reaction mixture was purified by reversed phase chromatography to give *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-4-(2,5-dimethylphenyl)-4-oxo-butyramide trifluoroacetate salt as a colorless oil (33 mg).

5

Other examples:

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxo-butyric acid with (4-acetylphenoxy)acetic acid gave 2-(4-acetylphenoxy)-*N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]acetamide.

10

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxo-butyric acid with *N*-phenylsulfonylglycine gave 2-benzenesulfonylamino-*N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]acetamide.

15

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(6-methoxynaphth-2-yl)-2-methylacetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(6-methoxynaphth-2-yl)propionamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 3-benzenesulfonylpropionic acid gave 3-benzenesulfonyl-*N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]propionamide.

20

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with (4-thiophen-2-ylpyrazol-1-yl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(4-thiophen-2-ylpyrazol-1-yl)acetamide.

25

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(chloro-3-methylbenzo[b]thiophen-2-yl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(chloro-3-methylbenzo[b]thiophen-2-yl)-acetamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(4-benzyloxyphenoxy)acetic acid gave 2-(4-benzyloxyphenoxy)-*N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]acetamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-methyl-2-(4-thiophenoylphenyl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(4-thiophenoylphenyl)propionamide.

5 Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(1-acetylnaphth-2-yloxy)-2-methylacetic acid gave 2-(1-acetylnaphth-2-yloxy)-*N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]propionamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with *N*-benzoylglycine gave 2-benzoylamino-*N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]acetamide.

10 Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(5,6-dimethylbenzimidazol-1-yl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(5,6-dimethylbenzimidazol-1-yl)acetamide.

15 Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(5-methyl-2-phenyloxazol-3-yl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(5-methyl-2-phenyloxazol-3-yl)acetamide.

20 Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(3-methyl-2-*N*-phenylpyrazol-4-yl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(3-methyl-2-*N*-phenylpyrazol-4-yl)acetamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(2-pyrazin-2-ylthiazol-4-yl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(2-pyrazin-2-ylthiazol-4-yl)acetamide.

25 Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(4-methyl-[1,2,3]thiadiazol-5-ylsulfanyl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(4-methyl-[1,2,3]thiadiazol-5-ylsulfanyl)-acetamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(naphth-2-ylsulfanyl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(naphth-2-ylsulfanyl)acetamide.

5 Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(quinoxalin-2-ylsulfanyl)acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(quinoxalin-2-ylsulfanyl)acetamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(2-chloro-4-fluorophenylsulfanyl)acetic acid gave 2-(2-chloro-4-fluorophenylsulfanyl)-*N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]acetamide.

10 Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-[2-(pyridin-2-yl)-6-trifluoromethylpyrimidin-4-ylsulfanyl]acetic acid gave *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-[2-(pyridin-2-yl)-6-trifluoromethylpyrimidin-4-ylsulfanyl]acetamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-[5-(4-chlorophenyl)pyrimidin-4-ylsulfanyl]acetic acid gave 2-[5-(4-chlorophenyl)pyrimidin-4-yl-sulfanyl]-*N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]acetamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(3,4-methylenedioxyphenyl)acetic acid and 3-aminomethyl-1-(3,4-dichlorobenzyl)pyrrolidine with 3-aminomethyl-1-(3,4-methylenedioxybenzyl)pyrrolidine gave *N*-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(3,4-methylenedioxy-phenyl)acetamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(3-phenylpyrazol-1-yl)acetic acid and 3-aminomethyl-1-(3,4-dichlorobenzyl)pyrrolidine with 3-aminomethyl-1-benzylpyrrolidine gave *N*-[1-(benzyl)-pyrrolidin-3-(*RS*)-ylmethyl]-2-(3-phenylpyrazol-1-yl)acetamide.


Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(3-fluoro-4-hydroxyphenyl)acetic acid and 3-aminomethyl-1-(3,4-dichlorobenzyl)pyrrolidine with 3-(*S*)-aminomethyl-1-(2,3-dichlorobenzyl)-

pyrrolidine gave *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*S*)-ylmethyl]-2-(3-fluoro-4-hydroxyphenyl)-acetamide.

Proceeding as described above but substituting 4-(2,5-dimethylphenyl)-4-oxobutyric acid with 2-(3-acetylaminophenyl)acetic acid and 3-aminomethyl-1-(3,4-dichlorobenzyl)pyrrolidine, with 3-(*S*)-aminomethyl-1-(2,3-dichlorobenzyl)pyrrolidine gave *N*-[1-(2,3-dichlorobenzyl)pyrrolidin-3-(*S*)-ylmethyl]-2-(3-acetylaminophenyl)-acetamide.

Example 11

Synthesis of 1-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-*m*-tolylurea TFA salt



Neat 3-methylphenylisocyanate (16 mg, 0.12 mmol, 1.1 equiv.) was added to a solution of 3-(*RS*)-aminomethyl-1-(3,4-methylenedioxybenzyl)pyrrolidine (28 mg, 0.12 mmol) in tetrahydrofuran (1 mL) at room temperature. The reaction mixture was shaken overnight, concentrated to dryness and diluted with dimethyl sulfoxide/methanol (1:1, 1 mL). The solution was purified using reversed phase chromatography to give, after evaporation of the solvents, 1-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-*m*-tolylurea as the TFA salt (26 mg).

Other examples:

Proceeding as described above but substituting 3-methylphenylisocyanate with 4-methoxyphenylisocyanate gave 1-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-(4-methoxyphenyl)urea.


Proceeding as described above but substituting 3-methylphenylisocyanate with phenethylisocyanate gave 1-[1-(3,4-methylenedioxybenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-(2-phenylethyl)urea.

Proceeding as described above but 3-aminomethyl-1-(3,4-methylenedioxybenzyl)pyrrolidine with 3-(*RS*)-aminomethyl-1-(3-phenylbenzyl)pyrrolidine gave 1-[1-(3-phenylbenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-*m*-tolylurea.

Proceeding as described above but substituting 3-(*RS*)-aminomethyl-1-(3,4-methylene-dioxybenzyl)pyrrolidine with 3-aminomethyl-1-(3,4-dichlorobenzyl)pyrrolidine and 3-methylphenylisocyanate with 1-(naphth-1-yl)ethylisocyanate gave 1-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-3-[1-(naphth-1-yl)ethyl]urea.

Example 12

Synthesis of *N*-[1-(3,4-dichlorobenzyl)pyrrolidin-3-(*RS*)-ylmethyl]-2-(9H-1,3,4,9-tetraazafluoren-2-ylsulfanyl)acetamide.



15

Solid (9H-1,3,4,9-tetraazafluoren-2-ylsulfanyl)acetic acid (390 mg, 1.5 mmol, 1.2 equiv.) was added to a solution of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (360 mg, 1.9 mmol, 1.5 equiv.), 1-hydroxybenzotriazole hydrate (260 mg, 1.9 mmol, 1.5 equiv.) and triethylamine (0.65 mL, 4.6 mmol, 3.7 equiv.) in chloroform (10 mL). 3-(*RS*)-Aminomethyl-1-(3,4-dichlorobenzyl)pyrrolidine (320 mg, 1.2 mmol) was added and the suspension was made homogeneous upon addition of dimethylformamide (5 mL). The brown solution was stirred for 14 h, then concentrated *in vacuo*. The residue was diluted with ethyl acetate and washed with 1 M sodium hydroxide. The organic phase was concentrated to dryness and the residue purified by chromatography (5→10% methanol in CHCl₃) to provide *N*-[1-(3,4-

25

dichlorobenzyl)-pyrrolidin-3-(*RS*)-ylmethyl]-2-(9H-1,3,4,9-tetraazafluoren-2-ylsulfanyl)acetamide as a pale yellow solid. The solid was dissolved into 4 M HCl in dioxane (1 mL) and the mixture concentrated under vacuum to afford pure product as its hydrochloride salt (250 mg).

Example 13

Formulation Examples

The following are representative pharmaceutical formulations containing a compound of Formula (I).

Tablet formulation

The following ingredients are mixed intimately and pressed into single scored tablets.

Ingredient	Quantity per tablet, mg
compound of this invention	400
cornstarch	50
croscarmellose sodium	25
lactose	120
magnesium stearate	5

Capsule formulation

The following ingredients are mixed intimately and loaded into a hard-shell gelatin capsule.

Ingredient	Quantity per capsule, mg
compound of this invention	200
lactose, spray-dried	148
magnesium stearate	2

Suspension formulation

The following ingredients are mixed to form a suspension for oral administration.

	Ingredient	Amount
5	compound of this invention	1.0 g
	fumaric acid	0.5 g
	sodium chloride	2.0 g
	methyl paraben	0.15 g
	propyl paraben	0.05 g
10	granulated sugar	25.5 g
	sorbitol (70% solution)	12.85 g
	Veegum K (Vanderbilt Co.)	1.0 g
	flavoring	0.035 ml
	colorings	0.5 mg
15	distilled water	q.s. to 100 ml

Injectable formulation

The following ingredients are mixed to form an injectable formulation.

	Ingredient	Amount
20	compound of this invention	0.2 g
	sodium acetate buffer solution, 0.4 M	2.0 ml
	HCl (1N) or NaOH (1N)	q.s. to suitable pH
	water (distilled, sterile)	q.s. to 20 ml

25

Liposomal formulation

The following ingredients are mixed to form a liposomal formulation.

	Ingredient	Amount
5	compound of this invention	10 mg.
	L- α -phosphatidylcholine	150 mg
	<i>tert</i> -butanol	4 ml

Freeze dry the sample and lyophilize overnight. Reconstitute the sample with 1 ml
10 0.9% saline solution. Liposome size can be reduced by sonication

Example 14CCR-3 Receptor Binding Assay- *in vitro*

15 The CCR-3 antagonistic activity of the compounds of the invention was determined by their ability to inhibit the binding of 125 I cotaxin to CCR-3 L1.2 transfectant cells ((see Ponath, P.D. et al., *J. Exp. Med.*, Vol. 183, 2437-2448, (1996)).

The assay was performed in Costar 96-well polypropylene round bottom plates.
20 Test compounds were dissolved in DMSO and then diluted with binding buffer (50 mM HEPES, 1 mM CaCl_2 , 5 mM MgCl_2 , 0.5% bovine serum albumin (BSA), 0.02% sodium azide, pH 7.24) such that the final DMSO concentration was 2%. 25 μ l of the test solution or only buffer with DMSO (control samples) was added to each well, followed by the addition of 25 μ l of 125 I-eotaxin (100 pmol) (NEX314, New England
25 Nuclear, Boston, MA) and 1.5×10^5 of the CCR-3 L1.2 transfected cells in 25 μ l binding buffer. The final reaction volume was 75 μ l.

After incubating the reaction mixture for 1h at room temperature, the reaction was terminated by filtering the reaction mixture through polyethylenimine treated

Packard Unifilter GF/C filter plate (Packard, Chicago, IL.). The filters were washed four times with ice cold wash buffer containing 10 mM HEPES and 0.5M sodium chloride (pH 7.2) and dried at 65 °C for approximately 10 min.. 25 µl/well of Microscint-20™ scintillation fluid (Packard) was added and the radioactivity retained on the filters was determined by using the Packard TopCount™.

The IC₅₀ value (concentration of test compound required to reduce ¹²⁵I-eotaxin binding to the CCR-3 L1.2 transfected cells by 50%) for compounds in Tables I-V of the invention was between 0.02 and 200µM.

Example 15

Inhibition of Eotaxin mediated chemotaxis of CCR-3 L1.2 transfectant cells--

In vitro Assay

The CCR-3 antagonistic activity of the compounds of this invention was determined by measuring the inhibition of eotaxin mediated chemotaxis of the CCR-3 L1.2 transfectant cells, using a slight modification of the method described in Ponath, P. D. et al., *J. Clin. Invest.* 97: 604-612 (1996). The assay was performed in a 24-well chemotaxis plate (Costar Corp., Cambridge MA). CCR-3 L1.2 transfectant cells were grown in culture medium containing RPMI 1640, 10% Hyclone™ fetal calf serum, 55 mM 2-mercaptoethanol and Geneticin 418 (0.8 mg/ml). 18-24 hours before the assay, the transfected cells were treated with n-butyric acid at a final concentration of 5 mM/1 x 10⁶ cells/ml, isolated and resuspended at 1 x 10⁷ cells/ml in assay medium containing equal parts of RPMI 1640 and Medium 199 (M 199) with 0.5% bovine serum albumin.

Human eotaxin suspended in phosphate buffered saline at 1 mg/ml was added to bottom chamber in a final concentration of 100 nM. Transwell culture inserts (Costar Corp., Cambridge MA) having 3 micron pore size were inserted into each well and L1.2 cells (1 x 10⁶) were added to the top chamber in a final volume of 100µl. Test

compounds in DMSO were added both to the top and bottom chambers such that the final DMSO volume was 0.5%. The assay was performed against two sets of controls. The positive control contained cells with no test compound in the top chamber and only eotaxin in the lower chamber. The negative control contained cells with no test compound in the top chamber and neither eotaxin nor test compound in lower
5 chamber. The plate was incubated at 37 °C. After 4 h, the inserts were removed from the chambers and the cells that had migrated to the bottom chamber were counted by pipetting out 500 µl of the cell suspension from the lower chamber to 1.2 ml Cluster tubes (Costar) and counting them on a FACS for 30 sec.

10

The IC₅₀ value (concentration of test compound required to reduce eotaxin mediated chemotaxis of CCR-3 L 1.2 transfected cells by 50%) for representative compounds of the invention was between 0.006 to 1.1 µM.

15

Example 16

Inhibition of Eotaxin mediated chemotaxis of human eosinophils--

In vitro Assay

20

The ability of compounds of the invention to inhibit eotaxin mediated chemotaxis of human eosinophils was assessed using a slight modification of procedure described in Carr, M.W. et al., *Proc. Natl. Acad. Sci. USA*, 91: 3652-3656 (1994). Experiments were performed using 24 well chemotaxis plates (Costar Corp., Cambridge MA). Eosinophils were isolated from blood using the procedure described in PCT Application, Publication No. WO 96/22371. The endothelial cells used were
25 the endothelial cell line ECV 304 obtained from European Collection of Animal Cell Cultures (Porton Down, Salisbury, U.K.). Endothelial cells were cultured on 6.5 mm diameter Biocoat® Transwell tissue culture inserts (Costar Corp., Cambridge MA) with a 3.0 µM pore size. Culture media for ECV 304 cells consisted of M199, 10% Fetal Calf Serum, L-glutamine and antibiotics. Assay media consisted of equal parts

RPMI 1640 and M199, with 0.5% BSA. 24 h before the assay 2×10^5 ECV 304 cells were plated on each insert of the 24-well chemotaxis plate and incubated at 37 °C. 20 nM of eotaxin diluted in assay medium was added to the bottom chamber. The final volume in bottom chamber was 600 µl. The endothelial coated tissue culture inserts were inserted into each well. 10⁶ eosinophil cells suspended in 100 µl assay buffer were added to the top chamber. Test compounds dissolved in DMSO were added to both top and bottom chambers such that the final DMSO volume in each well was 0.5%. The assay was performed against two sets of controls. The positive control contained cells in the top chamber and eotaxin in the lower chamber. The negative control contained cells in the top chamber and only assay buffer in the lower chamber. The plates were incubated at 37 °C in 5% CO₂/95% air for 1-1.5 h.

The cells that had migrated to the bottom chamber were counted using flow cytometry. 500 µl of the cell suspension from the lower chamber was placed in a tube, and relative cell counts were obtained by acquiring events for a set time period of 30 seconds.

Example 17

Inhibition of Eosinophil influx into the lungs of Ovalbumin sensitized balb/c mice by CCR-3 Antagonist --*In vivo* Assay


The ability of the compounds of the invention to inhibit leukocyte infiltration into the lungs was determined by measuring the inhibition of eosinophil accumulation into the bronchioalveolar lavage (BAL) fluid of Ovalbumin (OA)-sensitized balb/c mice after antigen challenge by aerosol. Briefly, male balb/c mice weighing 20-25g were sensitized with OA (10 µg in 0.2 ml aluminum hydroxide solution) intraperitoneally on days 1 and 14. After a week, the mice were divided into ten groups. Test compound or only vehicle (control group) or anti-eotaxin antibody (positive control group) was administered either intraperitoneally, subcutaneously or

orally. After 1 h, the mice were placed in a Plexiglass box and exposed to OA aerosol generated by a PARISTAR™ nebulizer (PARI, Richmond, VA) for 20 min. Mice which had not been sensitized or challenged were included as negative control. After 24 or 72 h, the mice were anesthetized (urethane, approx. 1 g/kg, i.p.), a tracheal cannula (PE 60 tubing) was inserted and the lungs were lavaged four times with 0.3 ml PBS. The BAL fluid was transferred into plastic tubes and kept on ice. Total leukocytes in a 20 µl aliquot of the BAL fluid was determined by Coulter Counter™ (Coulter, Miami, FL). Differential leukocyte counts were made on Cytospin™ preparations which had been stained with a modified Wright's stain (Diff-Quick™) by light microscopy using standard morphological criteria.

The ID₅₀ for the compounds of this invention in this assay is between 30 and 50 mgs/kg.

Claims

1. A compound selected from compounds of Formula (I):



Z is -N- or -(N⁺R)-, X⁻ wherein R is alkyl, haloalkyl, aralkyl, hydroxyalkyl, carboxyalkyl, alkoxyacetylalkyl, or cyanoalkyl, and X⁻ is a pharmaceutically acceptable counterion;

Ar¹ and Ar² are, independently of each other, aryl or heteroaryl;

Q is straight or branched alkylene with 1-3 carbon atoms;

R¹ is hydrogen or alkyl;

A is either:

(I) -N(R²)C(O)- when:

B is:

- (i) alkylene with 1-4 carbon atoms inclusive wherein one of the carbon atoms may optionally be replaced by a group selected from -C(O)-, -N(R⁴)-, -O-, -S(O)_n- (where n is 0, 1 or 2), -NR⁵C(O)- and -N(R⁶)SO₂-; or

(ii) an alkynylene chain;

wherein:

R² is hydrogen, alkyl, acyl, haloalkyl, heteroalkyl,

heterocyclalkyl, or -(alkylene)-C(O)-Z' where Z' is alkyl, haloalkyl,

alkoxy, haloalkyloxy, hydroxy, amino, or mono- or disubstituted amino;
and

R^4 , R^5 and R^6 are, independently of each other, hydrogen, alkyl,
acyl, haloalkyl, heteroalkyl, heterocyclalkyl, or $-(\text{alkylene})-\text{C}(\text{O})-\text{Z}'$
where Z' is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, or
mono- or disubstituted amino; or

(II) a group selected from $-\text{N}(\text{R}^2)\text{C}(\text{S})-$, $-\text{N}(\text{R}^2)\text{C}(\text{O})\text{N}(\text{R}^3)-$,
 $-\text{N}(\text{R}^2)\text{C}(\text{S})\text{N}(\text{R}^3)-$, $-\text{N}(\text{R}^2)\text{SO}_2-$, $-\text{N}(\text{R}^2)\text{SO}_2\text{N}(\text{R}^3)-$,
 $-\text{N}(\text{R}^2)\text{C}(\text{O})\text{O}-$, and $-\text{OC}(\text{O})\text{N}(\text{R}^3)-$ when:

B is:

- (i) a bond;
- (ii) an alkylene chain of 1-4 carbon atoms inclusive wherein one of
the carbon atoms may optionally be replaced by a group selected
from $-\text{C}(\text{O})-$, $-\text{N}(\text{R}^4)-$, $-\text{O}-$, $-\text{S}(\text{O})_n-$ (where n is 0, 1 or 2),
 $-\text{NR}^5\text{C}(\text{O})-$ and $-\text{N}(\text{R}^6)\text{SO}_2-$;
- (iii) an alkenylene chain; or
- (iv) an alkynylene chain;

wherein:

R^3 is hydrogen, alkyl, acyl, haloalkyl, heteroalkyl,
heterocyclalkyl, or $-(\text{alkylene})-\text{C}(\text{O})-\text{Z}'$ where Z' is alkyl, haloalkyl,
alkoxy, haloalkyloxy, hydroxy, amino, or mono- or disubstituted amino;
and

R^2 , R^4 , R^5 and R^6 are as defined above; and

prodrugs, individual isomers, mixtures of isomers and pharmaceutically acceptable
salts thereof.

2. The compound of Claim 1 wherein Z is $-\text{N}-$; R^1 is hydrogen and A is
 $-\text{NHC}(\text{O})-$.

3. The compound of Claim 2 wherein Ar¹ is a naphthyl or a phenyl ring substituted with one, two, or three substituents selected from alkyl, cyano, nitro, halo, methylenedioxy, ethylenedioxy, alkoxy or phenoxy.
- 5 4. The compound of Claim 3 wherein Q and B are -CH₂-.
5. The compound of Claim 4 wherein Ar² is an aryl ring.
6. The compound of Claim 5 wherein:
10 Ar¹ is a phenyl ring substituted with one, two, or three substituents selected from methyl, chloro, fluoro, bromo, or methylenedioxy; and
Ar² is a phenyl ring optionally substituted with one, two, or three substituents selected from alkoxy, alkylthio, halo, amino, -NHC(O)R' (where R' is alkyl or optionally substituted phenyl), hydroxy, or -SO₂Me.
- 15 7. The compound of Claim 3 wherein:
Q is -CH₂- and B is -(CH₂)₂;
8. The compound of Claim 7 wherein Ar² is a heteroaryl ring.
- 20 9. The compound of Claim 8 wherein:
Ar¹ is a phenyl ring substituted with one, two, or three substituents methyl, chloro, fluoro, bromo, or methylenedioxy.
- 25 10. The compound of Claim 9 wherein Ar¹ is 3-chlorophenyl, 4-chlorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-ethylenedioxyphenyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, or 3,4-methylenedioxyphenyl.


11. The compound of Claim 10 wherein Ar^2 is a pyrimidin-2-yl ring optionally substituted at the 5-position with a phenyl ring which is optionally substituted with one, two, or three substituents selected from methyl, methoxy, methylthio, chloro, fluoro, vinyl, or dimethylamino.
- 5 12. The compound of Claim 3 wherein Q is $-\text{CH}_2-$ and B is $-\text{CH}_2\text{S}-$.
13. The compound of Claim 12 wherein:
 Ar^2 is a heteroaryl ring.
- 10 14. The compound of Claim 13 wherein:
 Ar^1 is a phenyl ring optionally substituted with one, two, or three substituents methyl, chloro, fluoro, bromo, or methylenedioxy.
- 15 15. The compound of Claim 14 wherein Ar^1 is 3-chlorophenyl, 4-chlorophenyl, 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-ethylenedioxyphenyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, or 3,4-methylenedioxyphenyl.
- 20 16. The compound of Claim 15 wherein Ar^2 is a pyrimidin-2-yl ring optionally substituted at the 5-position with a phenyl ring which optionally substituted with one, two, or three substituents selected from methyl, methoxy, methylthio, chloro, fluoro, vinyl or dimethylamino.
- 25 17. The compound of Claim 3 wherein Q is $-\text{CH}_2-$ and B is $-\text{CH}_2\text{O}-$.
18. The compound of Claim 17 wherein:
 Ar^2 is a heteroaryl ring.
- 30 19. The compound of Claim 18 wherein:

Ar¹ is a phenyl ring optionally substituted with one, two, or three substituents methyl, chloro, fluoro, bromo, or methylenedioxy.

20. The compound of Claim 19 wherein Ar¹ is 3-chlorophenyl; 4-chlorophenyl,
5 3,4-difluorophenyl, 3-chloro-4-fluorophenyl, 4-chloro-3-fluorophenyl, 4-fluoro-3-methoxyphenyl, 2,4-difluorophenyl, 3,4-ethylenedioxyphenyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, or 3,4-methylenedioxyphenyl.
21. The compound of Claim 20 wherein Ar² is a pyrimidin-2-yl ring optionally
10 substituted at the 5-position with a phenyl ring which optionally substituted with one, two, or three substituents selected from methyl, methoxy, methylthio, chloro, fluoro, vinyl or dimethylamino.
22. The compound of Claim 1 wherein R¹ is hydrogen and A is -NHC(O)NH-.
15
23. The compound of Claim 22 wherein Ar¹ is a naphthyl or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, cyano, nitro, halo, methylenedioxy, ethylenedioxy, alkoxy or phenoxy.
- 20 24. The compound of Claim 23 wherein Q is -CH₂- and B is a bond or -CH₂-.
25. The compound of Claim 24 wherein Ar² is an aryl ring.
26. The compound of Claim 25 wherein:
25 Ar¹ is a phenyl ring optionally substituted with one, two, or three substituents selected from methyl, chloro, fluoro, bromo, or methylenedioxy; and
Ar² is a phenyl ring optionally substituted with one or two substituents selected from alkyl or alkoxy.
- 30

27. A pharmaceutical composition comprising a therapeutically effective amount of a compound of Claim 1 and a pharmaceutically acceptable excipient.


28. A process for preparing a compound of Claim 1, which comprises reacting a compound of Formula (IIa) where R^1 , Q, and Ar^1 are as defined in Claim 1:



- (i) with an acylating agent of formula $Ar^2-B-COL$ where L is a leaving group under acylating reaction conditions or an acid anhydride of formula $(Ar^2-B-CO)_2O$ to provide a compound of Formula (I) where A is $-N(R^2)CO-$ wherein R^2 is hydrogen; or
- (ii) with an amine of formula $Ar^2-B-NH(R^3)$ where R^3 is as defined in the Summary of the Invention, in the presence of a suitable coupling agent or an isocyanate of formula $Ar^2-B-N=C=O$ or a carbamoyl halide of formula $Ar^2-B-N(R^3)-C(O)L$ where R^3 is as defined in the Summary of the Invention and L is a halo group to provide a compound of Formula (I) where A is $-N(R^2)CON(R^3)-$ wherein R^2 is hydrogen; or
- (iii) with an amine of formula $Ar^2-B-NH(R^3)$ where R^3 is as defined in the Summary of the Invention, in the presence of a suitable coupling agent or an isothiocyanate of formula $Ar^2-B-N=C=S$ or a thiocarbamoyl halide of formula $Ar^2-B-N(R^3)-C(S)L$ where R^3 is as defined in the Summary of the Invention and L is a halo

- group to provide a compound of Formula (I) where A is $-N(R^2)C(S)N(R^3)-$ wherein R^2 is hydrogen; or
- (iv) with a sulfonylating agent of formula Ar^2-B-SO_2L or $Ar^2-B-N(R^3)-SO_2L$ where R^3 is as defined in the Summary of the Invention and L is a leaving group under sulfonylating reaction conditions to provide a compound of Formula (I) where A is $-N(R^2)SO_2-$ or $-N(R^2)SO_2N(R^3)-$ wherein R^2 is hydrogen; or
- (v) with an alcohol of formula Ar^2-B-OH in the presence of a suitable coupling agent to provide a compound of Formula (I) where A is $-N(R^2)C(O)O-$ wherein R^2 is hydrogen; and
- (vi) optionally converting a compound of Formula (I) prepared in Steps (i) to (v) above, where R^2 is hydrogen to a compound of Formula (I) where R^2 is not hydrogen; and
- (vii) optionally reacting a compound of Formula (I) prepared in Steps (i) to (vi) above, with an alkylating agent of formula RX where R is alkyl and X is a leaving group under alkylating conditions to provide a corresponding compound of Formula (I) where Z is $-(N^+R)-X^-$; and
- (viii) optionally replacing one counterion in the compound of Formula (I) prepared in Step (vii) above, with another counterion; and
- (ix) optionally converting the compound of Formula (I) prepared in Steps (i) to (viii) above, to the corresponding acid addition salt by treatment with an acid.

29. A process for preparing a compound of Claim 1, which comprises:



- (i) reacting a compound of formula (IIb) where R^1 , A, B and Ar^2 are as defined above, with an alkylating agent of formula Ar^1-Q-Y where Ar^1 is as defined in Claim 1 and Y is a leaving group under alkylating reaction conditions to provide a compound of Formula (I); and
- 5 (ii) optionally reacting a compound of Formula (I) prepared in Step (i) above, with an alkylating agent of formula RX where R is alkyl and X is a leaving group under alkylating conditions to provide a corresponding compound of Formula (I) where Z is $-(N^+R)-X^-$; and
- (iii) optionally replacing one counterion in the compound of Formula (I) prepared in
- 10 Step (ii) above, with another counterion; and
- (iv) optionally converting the compound of Formula (I) prepared in Steps (i) to (iii) above, to the corresponding acid addition salt by treatment with an acid.
30. A compound as claimed in claim 1 whenever prepared by a process as claimed
- 15 in claim 28 or 29.
31. A compound as claimed in any one of claims 1 to 26 as a therapeutic agent.
32. Use of a compound as claimed in any one of claims 1 to 26 for the preparation
- 20 of a medicament comprising as an active ingredient a compound as claimed in any one of claims 1 to 26 for the treatment of a disease treatable by administration of a CCR-3 antagonist, e.g. asthma.
33. A method of treatment of a disease in a mammal treatable by administration of
- 25 a CCR-3 antagonist, comprising administration to the mammal of a therapeutically effective amount of a compound of Claim 1.
34. The invention as hereinbefore described.

INTERNATIONAL SEARCH REPORT

In tional Application No
PCT/EP 99/08665

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D207/09 A61K31/40 C07D401/12 C07D401/14 C07D403/12
C07D403/14 C07D405/06 C07D405/12 C07D405/14 C07D409/06
C07D409/12 C07D409/14 C07D487/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	EP 0 916 668 A (BANYU PHARMA CO LTD) 19 May 1999 (1999-05-19) the whole document	1-34
A	& WO 98 04554 A (BANYU PHARMA CO LTD) 5 February 1998 (1998-02-05) the whole document	1-34
A	US 3 459 757 A (BRABANDER HERBERT JOSEPH ET AL) 5 August 1969 (1969-08-05) the whole document	1-34
P, X	EP 0 903 349 A (HOFFMANN LA ROCHE) 24 March 1999 (1999-03-24) page 64 -page 65; claim 1	1-11, 22-28, 30-34

-/-



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

23 February 2000

Date of mailing of the international search report

19.03.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Fink, D

INTERNATIONAL SEARCH REPORT

In International Application No
PCT/EP 99/08665

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>WO 99 25686 A (TSUTSUMI TAKAHARU.; FURUYA MONORU (JP); HADA TAKAHIKO (JP); IMAI MI) 27 May 1999 (1999-05-27)</p> <p>page 355 -page 357; claim 1</p> <p>page 53; table 1.33, compound no. 353</p> <p>pages 153-154; tables 1.133 and 1.134, compounds no. 1463-1471</p> <p>pages 157-158; tables 1.137 and 1.138, compounds no. 1506-1513</p>	1-3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 99/08665

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 33 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 1-34 (partly)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-34 (partly)

Present claims 1-26 inter alia relate to prodrugs of the compounds of the present formula (I). The term prodrug in the present context is considered to lead to a lack of clarity within the meaning of Article 6 PCT (this term does not provide any information in respect of the structure of these compounds). It is thus impossible to compare the full scope of the present compound claims with what is set out in the prior art. The lack of clarity is such as to render a meaningful complete search impossible. Consequently, the search has been restricted to the compounds of the present formula (I) as defined in the present claim 1 (including their individual isomers, mixtures of their isomers and pharmaceutically acceptable salts thereof), pharmaceutical compositions containing them, processes for preparing them, and their use (i.e., the International Search Report is incomplete with respect to the prodrugs of the present compounds).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/08665

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0916668 A	19-05-1999	AU 3633997 A WO 9804554 A	20-02-1998 05-02-1998
US 3459757 A	05-08-1969	BE 663869 A BE 687242 A BE 688357 A BE 688358 A CH 473133 A CH 497436 A CH 502356 A DE 1568414 A DE 1695161 A DE 1695162 A FR 4401 M FR 7192 M FR 5905 M FR 1495674 A FR 1499512 A FR 1502968 A GB 1083724 A GB 1124157 A GB 1135336 A GB 1156042 A NL 6613466 A NL 6614923 A NL 6614924 A US 3299044 A US 3305560 A US 3317538 A US 3393208 A US 3426046 A US 3429885 A US 3462462 A US 3481921 A US 3560352 A DK 120548 B DK 111489 B IL 26724 A IL 26725 A NO 118326 B NO 118918 B SE 334620 B SE 329856 B	16-11-1965 22-03-1967 17-04-1967 17-04-1967 31-05-1969 15-10-1970 31-01-1971 02-04-1970 10-12-1970 10-12-1970 18-08-1969 25-03-1968 20-12-1967 17-01-1968 07-02-1968 25-06-1969 28-03-1967 24-04-1967 24-04-1967 17-01-1967 21-02-1967 02-05-1967 16-07-1968 04-02-1969 25-02-1969 19-08-1969 02-12-1969 02-02-1971 14-06-1971 02-09-1968 30-11-1970 17-09-1970 15-12-1969 02-03-1970 03-05-1971 26-10-1970
EP 0903349 A	24-03-1999	AU 8080098 A CA 2245043 A CN 1211572 A CZ 9802566 A DE 19837386 A FR 2767826 A GB 2330580 A HR 980450 A HU 9801887 A JP 11147872 A NO 983749 A PL 328049 A	25-02-1999 18-02-1999 24-03-1999 17-03-1999 25-02-1999 05-03-1999 28-04-1999 30-06-1999 28-06-1999 02-06-1999 19-02-1999 01-03-1999
WO 9925686 A	27-05-1999	AU 1374199 A	07-06-1999

THIS PAGE BLANK (USPTO)